

Evolution of ancient harbours in deltaic contexts: A geoarchaeological typology

Matthieu Giaime^{a,*}, Nick Marriner^b, Christophe Morhange^c

^a Department of Geography, Durham University, South Road, Durham DH1 3LE, UK

^b CNRS, Laboratoire Chrono-Environnement UMR 6249, Université de Franche-Comté, UFRST, 16 Route de Gray, 25030 Besançon, France

^c Aix Marseille Univ, CNRS, IRD, INRA, Coll France, CEREGE UMR 7330, Europôle Méditerranéen de l'Arbois, BP80, 13545 Aix-en-Provence, France



ARTICLE INFO

Keywords:

Delta
Ancient harbour
Sediment supply
Lagoon
Estuary
Landscape archaeology
Geoarchaeology
Geomorphology
Coasts
Relative sea-level changes
Mediterranean
Black Sea
Holocene

ABSTRACT

In deltaic areas, within a context of relative sea-level stability during the past 6000 years, environmental conditions have been key in mediating human settlements and the evolution of ancient harbours. The natural diversity of clastic-coast environments, in particular deltas, is central to explain the wide disparity in harbour contexts. It is, therefore, important to understand the impact of these settlements on the evolution of clastic coasts. How did ancient societies exploit and adapt to deltaic environments? Here, we detail a typology of ancient harbours on clastic coasts using a suite of multidisciplinary case studies from the Mediterranean and the Black Sea.

We present the impact of different forcing agents (floods, storms, sedimentary inputs, relative sea-level (RSL) changes, dredging and harbour structures) on the geomorphological evolution of selected harbour sites, and underline the important role of coastal changes at different temporal and spatial scales. These processes affected the infrastructure and the viability of harbours to varying degrees. In deltaic contexts, our analysis differentiates five harbour types: (1) fluvial harbours, affected by floods, rapid sedimentation and river-channel changes; (2a) infilled estuarine harbours; or (2b) submerged estuarine harbours, lying at the interface between marine and fluvial processes; (3) lagoonal harbours strongly affected by sedimentary inputs; and (4) the impact of humans on harbour basins via harbour protection structures and dredging. For these harbour types, we probe the advantages and disadvantages of each geomorphological setting, in addition to the consequences of environmental pressures on human societies.

1. Introduction

The Holocene morphogenesis of Mediterranean deltas can be divided into two main successive steps (Anthony et al., 2014). First, the glacio-eustatic stabilisation of sea level since ca. 6000 years, associated with continental sedimentary inputs, favoured the widespread progradation of clastic coasts (Stanley and Warne, 1994, 1997; Hori et al., 2004; Catuneanu and Zecchin, 2013; Anthony et al., 2014). Subsequently, the importance of human impacts, mainly linked to watershed deforestation, led to soil erosion and an increase in fluvial sediment supply that accentuated the expansion of deltaic lobes until the mid-20th century (e.g. Bellotti et al., 2007; Simeoni and Corbau, 2009; Maselli and Trincardi, 2013).

Deltaic environments, which constitute depocenters for fluvial sediments, have been occupied by human societies since the stabilisation

of sea level during the mid-Holocene acting, in many instances, as cradles for the emergence of early social complexity (Stanley and Warne, 1997; Day et al., 2007; Kennett and Kennett, 2006). In the Western Mediterranean, human influence has been highlighted in the formation of several Italian deltas. Marchetti (2002) proposed that human impacts have been decisive in modulating the evolution of the Po River since Roman times due to the intensive exploitation of its catchment between the 1st and the 3rd centuries AD. More locally, the deltas of the rivers Arno and Ombrone manifest a similar evolution and chronology. Their morphogenesis is mostly attributed to the past 2500 years, following the massive deforestation of their catchments (Pranzini, 2001, 2007; Bellotti et al., 2004; Amorosi et al., 2013). This contemporaneous evolution of deltas during the last 2500 years has led to the term “man-made deltas” being coined (Maselli and Trincardi, 2013).

* Corresponding author.

E-mail addresses: matthieu.giaime@gmail.com, matthieu.j.giaime@durham.ac.uk (M. Giaime), nick.marriner@univ-fcomte.fr (N. Marriner), morhange@cerege.fr (C. Morhange).

<https://doi.org/10.1016/j.earscirev.2019.01.022>

Received 3 July 2018; Received in revised form 15 January 2019; Accepted 23 January 2019

Available online 25 January 2019

0012-8252/ © 2019 Elsevier B.V. All rights reserved.




Fig. 1. Location of the sites (harbours or deltas) discussed in this manuscript.

Progradation led to important changes in coastal landscapes and had a direct impact on littoral societies leading, in many well-documented cases, to the deterioration of harbour accessibility and functioning (see Fig. 1 for the location of the sites mentioned in this paper). In Asia Minor, Miletus' and Ephesus' ancient harbours, located in ria-type deltaic environments are now located several tens of kilometres inland and underwent important environmental changes over time (Brückner et al., 2017). These ria-type deltas evolved following a three-phase model (Anthony et al., 2014). At Ephesus, Brückner (2005), Stock et al. (2013, 2014, 2016) and Brückner et al. (2006), have shown that these three phases comprised: (i) the flooding of a tectonic graben by postglacial sea-level rise, creating a fluvial embayment around 8000 to 6000 years ago. Such environments were particularly conducive to human settlement because they lay in a sheltered position compared to the open coast and they had direct access to freshwater due to the proximity of the river. (ii) The rapid westward progradation of the deltaic front of the Küçük Menderes since 6000–5000 years BP, much like the nearby ria of the Büyük Menderes, illustrates the first environmental pressures that can affect harbours (Brückner et al., 2014, 2006); and (iii) a straightening of the shoreline at the ria mouth, engendering a landlocking of the harbours on the deltaic plain. In the same vein, geoarchaeological research on the Acheloos delta in Greece (Epirus) has highlighted the effects of deltaic progradation on shoreline changes and harbour landlocking (Vött, 2007; Vött et al., 2007; Fig. 2). The shore of the palaeo-island of Triardo, which accommodated the city of Oiniadai, was washed by the sea 6000 years ago because of sea-level rise. Thereafter, the progradation of the Acheloos River delta isolated the island from the sea. The ancient harbours remained connected to the sea via fluvial arms, which acted as communication channels. By Roman and Byzantine times, the island was permanently landlocked and integrated into the deltaic plain. This limitation was overcome by the establishment of a fluvial harbour, on the bank of a meander of the Acheloos, on the southern part of the island. This relocation of the harbour is typical of the “race to the sea” induced by the environmental changes at that time.

Deltaic environments show specific geomorphological and ecological particularities, and offered numerous natural potentialities for human societies and harbour activities during Antiquity. The objective of this paper is to demonstrate that environmental changes have played

a major role in the geomorphological evolution of ancient harbours in deltaic contexts since the Bronze Age. We also shed light on the potentialities that deltaic environments offered for societies and commercial activities, as well as the role of human impacts in shaping the evolution of harbours. For this purpose, we use case-study examples to probe the impacts of the diverse environmental pressures on the evolution of deltaic harbours. We highlight the (dis)advantages of each geomorphological context identified.

2. Defining a typology

The diversity of deltaic environments is illustrated by the wide plethora of geomorphological contexts in which harbours were located. It is interesting to define a geoarchaeological typology that translates the geomorphological trajectory of ancient deltaic harbours over time. The typology includes the influence of rivers, marine processes and human societies. We identified several pressures that affect harbours with respect to their geomorphological context (mainly their location in relation to the river channel/mouth or to the sea).

2.1. Identification of the forcing agents

We detail different environmental pressures and human impacts. Because deltas are depocenters for sediments eroded from the watershed, one key environmental variable affecting harbours over time is fluvial sediment supply. Over the last 6000 years, Mediterranean clastic coasts have undergone an important progradation at all spatial scales, after the deceleration of sea-level rise and an increase in fluvial sediment supply to base-level depocenters (Stanley and Warne, 1994; Stewart and Morhange, 2009; Syvitski and Kettner, 2011; Maselli and Trincardi, 2013; Anthony et al., 2014). These factors explain why ancient populations had to constantly adapt to coastal progradation and landscape change. In Cyprus, Devillers (2008) demonstrated that the Gialias ria was flooded by the sea 8000 years ago, following the post-glacial sea-level rise. He elucidated a succession of coastal palaeoenvironments (alluvial plains, lagoons, coastal spits) consistent with the eastward migration of the coastline during the past 6000 years. Since the Bronze Age, local populations have had to adapt to these environmental changes and relocate their harbours to keep pace with




Fig. 2. A. Palaeogeography of the progradation of the Acheloos delta since 600 BC and the location of the harbours of Oiniadai (Trikardo palaeo-island) mentioned in the text (adapted from Vött et al., 2007). B. Regional location of the site.




Fig. 3. A. Geomorphological map of the Gyalias ria (adapted from Devillers, 2008). B. Regional location of the site.

the shifting coastline. In Medieval times, this “race to the sea” led to the relocation of the harbour at Famagusta, on the waterfront outside of the infilled ria (Fig. 3).

At Utica (Mejerda delta, Tunisia) the importance of deltaic progradation in shaping harbour location and activities has been highlighted (Chelbi et al., 1995; Delile et al., 2015). Progradation of the delta from the south-west led to the development of a first deltaic lobe along the southern part of the promontory since the foundation of the Phoenician city (12th century BC), which hindered the development of maritime activities. The harbour had to be located north of the promontory and was linked to the sea via a maritime corridor west of the northern part of the delta (Chelbi et al., 1995; Delile et al., 2015).

Deltas are land-sea interfaces. We have identified two pressures linked to high-energy events. The first relates to river floods and the second to meteo-marine events affecting the coast. River floods can cause important destruction as attested at Pisa, where excavations in the harbour basin have led to the discovery of 16 ancient boats destroyed during flood events (Benvenuti et al., 2006; Allinne et al., 2016). At Narbonne (Sanchez et al., 2014; submitted; Faisse et al., 2018), repairs to the embankments of the artificialized channel of the Aude River were frequent during the Early Roman Empire and reveal the challenges of maintaining the river in its bed. These high-energy events raise the question of the impact of floods on the organisation of urban areas and ancient harbours, particularly in fluvial contexts (Arnaud-Fassetta et al., 2003; Benvenuti et al., 2006; Allinne, 2007; Arnaud-Fassetta et al., 2010; Leveau, 2012; Ruiz et al., 2014; Bini et al.,

2015; Allinne et al., 2016; Salomon et al., 2017).

Meteo-marine dynamics and high-energy events are well documented on clastic coastlines (Marriner et al., 2017a). In the Northern Adriatic Sea, Kaniewski et al. (2016) highlighted six different storm phases during the last 4500 years on the Mirna Delta (Croatia). These storm phases affected coastal agriculture and were correlated with periods of low solar activity, corroborating previous work by Camuffo et al. (2000). The impacts of storms are limited for naturally protected deltaic harbours (river mouths, lagoons), except in the case of exceptionally intense events, in contrast to open sea harbours located along the waterfront. Nonetheless, we would like to insist upon the fact that such evidence is difficult to observe in the sedimentary record. For instance, the sedimentological traits of storm facies are extremely challenging to differentiate from those generated by tsunamis (Dawson and Stewart, 2007; Morton et al., 2007; Engel and Brückner, 2011; Engel and May, 2012; Goff et al., 2012; Weiss, 2012; Pilarczyk et al., 2014; Marriner et al., 2017a).

The fourth forcing agent identified is related to RSL changes. During the Holocene, sea-level changes in tectonically stable regions are primarily linked to modifications in ice volumes and the Earth's response to the changing ice-water load. During the last 4000 years, the ice-equivalent melt-water input is considered minimal and does not exceed 1 m (Lambeck et al., 2004, 2014; Vacchi et al., 2016, 2018). In deltaic settings, the compaction of unconsolidated fine-grained sediments can lead to important RSL rise (Stanley, 1997; Törnqvist et al., 2008; Teatini et al., 2011; Marriner et al., 2012). In southern Italy, the Crati Delta,



Fig. 4. Dredging scars from the ancient harbour of Naples (Viviana Liuzza, Soprintenza archeologica di Napoli (A) and (B) Carsana et al., 2009). C. Leonardo da Vinci sketch depicting a dredging machine (draga cavafango; Manoscritto E, foglio 75 v.)

where the ancient Greek cities of Sybaris, Thuri and Copia are located, was affected by rapid progradation and long-term subsidence due to the combination of tectonics and sediment compaction (Pagliarulo, 2006; Stanley and Bernasconi, 2009, 2012; Cianflone et al., 2018). Tectonic activity can also play a role in RSL changes at short timescales. This variable, which can lead to important modifications in the shoreline and sometimes to the isolation of the port from the open sea, is nonetheless restricted to a small number of harbours, particularly in the central and eastern Mediterranean (Stewart and Morhange, 2009).

We identified human-related pressures highlighted by the variable density of harbour works. These reflect the evolution of harbour building techniques during Antiquity, and shed light on the importance of harbour dredging. Furthermore, at a wider geographic scale, human impacts played a key role in the intense deforestation of watersheds (Quézel and Médail, 2003) that, in conjunction with external dynamics (tectonics, climate variability) can lead to increases in fluvial sediment supply to coastal systems at base level (Dearing and Jones, 2003; Wainwright and Thornes, 2004; Casana, 2008; Syvitski and Kettner, 2011; Duser et al., 2011) and therefore the infilling of harbours.

Early coastal societies sought to reinforce natural coastal endowments in order to install permanent harbour areas (Blackman, 1982a, 1982b; Marriner et al., 2014, 2017b; Morhange et al., 2016b). The earliest evidence for artificial harbour works is dated to the Bronze Age and derives from the Red Sea. Archaeological surveys at Wadi al-Jafr (Egypt) have highlighted the presence of a large artificial anchorage, covering around 2.5 ha (0.02 km²), framed by an L-shaped mole dating to the Fourth Dynasty (2613–2494 BC; Tallet and Marouard, 2014, 2017). On clastic coasts, inhabitants favoured naturally-protected areas. But they could also manually dig basins, as was the case for Lechaion which was excavated out of a marsh area during the 6th century BC, at the mouth of a coastal stream (Stiros et al., 1996;

Morhange et al., 2012; Kolaiti et al., 2017). In Roman times (1st–2nd centuries BC), an intensification of harbour building, resulting from the discovery of hydraulic mortar (Pozzolana), allowed artificial basins to be erected on the seafront (Brandon et al., 2014). Several examples are known in the Mediterranean, such as Frejus (Gébara and Morhange, 2011) or *Portus*, with the construction of Claudius and Trajan basins (Keay et al., 2005; Goiran et al., 2010). The dredging of harbours is another example of human activities affecting harbour environments (Marriner and Morhange, 2006a; Morhange and Marriner, 2010). The most iconic example comes from the ancient harbour of Neapolis (Naples). The harbour bottom was completely reshaped by intensive dredging between the 4th and the 2nd centuries BC, leading to the absence of older sediments in the basin (Giampaola et al., 2004; Carsana et al., 2009; Di Donato et al., 2018; Fig. 4). The power of the dredging technology used is attested by the dimensions of the fossilised dredging scars, 80–165 cm wide and 30–50 cm deep. Such operations are recorded until the abandonment of the harbour in the 5th century AD (Di Donato et al., 2018).

2.2. Estimating the weight of forcing agents

We built a semi-quantitative database in which a percentage score was attributed to each forcing agent based on the scientific literature available for each site (see supplementary material for references). In Table 1, we detail the percentage values ascribed to the Bronze Age harbour of Tel Akko (Israel), to the Roman harbour of Ephesus (Turkey) and to the Eastern Harbour (Magnus Portus) of Alexandria (Egypt).

2.3. Statistical analyses

Based on these estimates, we created a heat map to visually

Table 1

Semi-quantified pressures for three harbours examples, Tel Akko (Israel), Ephesus (Turkey) and Magnus Portus of Alexandria (Egypt). The examples present three different harbours located in diverse geomorphological settings and for which the weight of each pressure is different. The different estimates for all the sites included in the present study are described in the Supplementary Material (Sivan et al., 2001).

Harbour	Period of use	Flood impacts	Sedimentary inputs	Storm impacts	Relative sea-level changes	Dredging	Infrastructure
Tel Akko Bronze and Iron Ages	2000 BC - 587 BC	Low	High	Low	Absent	Absent	Absent
		Possible low-energy floods	Infilling of the estuary attested	Marine-dominated estuary moderately exposed to the sea	Stable sea level since 4000 years	No direct evidence	No archaeological excavations
		∅	Morhange et al., 2016a; Giaime et al., 2018	Morhange et al., 2016a; Giaime et al., 2018	Sivan et al., 2001	∅	∅
Weight	2	3	2	0	0	0	
Percentage	28.5%	43%	28.5%	0%	0%	0%	
Harbour	Period of use	Flood impacts	Sedimentary inputs	Storm impacts	Relative sea-level changes	Dredging	Infrastructure
Ephesus	1 st c. BC – 7 th century AD	High	High	Possible	Possible	High	High
		River channel diverted under Hadrian (129 AD)	Rapid progradation of the delta front	Limited exposure to the open sea	Linked to global sea-level changes	Dredging of the harbour and excavation of the channel	Excavation of an artificial basin and channel
		Kraft et al., 2011; Stock et al., 2013, 2016	Stock et al., 2016	∅	∅	Stock et al., 2013, 2016	Kraft et al., 2011
Weight	4	4	1	1	4	4	
Percentage	22%	22%	6%	6%	22%	22%	
Harbour	Period of use	Flood impacts	Sedimentary inputs	Storm impacts	Relative sea-level changes	Dredging	Infrastructure
Alexandria -Eastern harbour (Magnus Portus)	4 th century BC – 7-8 th centuries AD	Absent	Moderate	Low	High	Possible	Moderate
		∅	Infilling of the harbours and presence of a submarine tomolo	High-energy event in the harbour but protected basin (presence of fine-grained sediments)	Subsidence of the delta front due to the compaction of unconsolidated sediments	No direct evidence	Submerged ancient structures in the Eastern harbour
		∅	Goiran et al., 2005; Marriner et al., 2008	Goiran et al., 2012	Stanley et al., 2006	∅	Goddio and Yoyotte, 2008; Goddio and Masson-Berghoff, 2016
Weight	0	3	2	4	1	3	
Percentage	0%	23%	15%	31%	8%	23%	




Fig. 5. Heat-map and associated cluster analysis of the ancient harbour dataset. The cluster analysis allowed us to distinguish four different groups (A to D) based on the importance of the environmental and/or anthropogenic forcings that affected the harbour basins. Details of the database are given in the Supplementary Material.

represent the data (Fig. 5). The semi-quantitative database was analysed using hierarchical clustering and Principal Components Analysis (PCA). We chose to apply multivariate statistical techniques because they facilitate the investigation of large and complex datasets, while taking into consideration the effects of all variables on the responses. In this study, cluster analysis (Algorithm: Paired group; Similarity measure: Correlation) was used to group data according to the estimated importance of each pressure identified on the harbour (Fig. 5). Subsequently, a PCA allowed us to test the ordination of the harbours according to the different pressures (Fig. 6).

The cluster analysis and the PCA highlight four different groups (noted A to D), based on the weight of the pressures affecting them (Figs. 5 and 6). In the PCA (Fig. 6), the main variance is loaded by the PCA-Axes 1 (49.17%) and 2 (25.15%). The groups highlighted by the cluster analysis are confirmed and are clearly differentiated by the PCA plot of the two main axes. PCA-Axis 1 contrasts harbours dominated by fluvial influences (positive scores) and harbours dominated by coastal/marine processes (negative scores). PCA Axis-2 characterizes the different harbour management strategies. Negative values are related to important human pressures on the shoreline in Antiquity while the harbours with modest human impacts are characterized by positive figures.

3. Presentation of the typology

The different forcing agents can be grouped according to their relationship to the river (floods and sedimentary inputs), the sea (storms and RSL changes) or human societies (dredging and harbour work). In this section, we present the different types of geomorphological contexts exploited by human societies for their harbours. We highlight the key potentialities of these contexts for harbour activities but also the main processes affecting the basins.

Rivers were, in ancient times, pivotal commercial and transport interfaces, between the coastline and the hinterland. Their key role explains why many important ancient cities were located in deltaic contexts, such as the important harbour network of Narbonne in connection with the Bages-Sigean lagoon. The city, founded in 118 BC at the crossroads between terrestrial (Via Domitia and Via Aquitania), maritime and fluvial (Aude/Garonne) axes was a pole of international trade. The harbour city attracted many ship owners and traders, and Narbonne was one of the most active harbours in the Roman Empire (Sanchez and Jezegou, 2011).

3.1. Fluvial harbours

The cluster analysis undertaken on the dataset includes the fluvial harbours (Arles, Pisa, Halmyris) in Group C, in addition to other




Fig. 6. Principal Components Analysis of the database. The figure presents the distribution of the harbour sites studied according to the importance of each environmental and/or anthropogenic pressure identified. The groups correspond to the cluster analysis results. The histograms denote the weight of each characteristic for Axis-1 and Axis-2.

harbours primarily subjected to flood hazards (e.g. river-mouth harbours). Fluvial harbours were generally located on riverbanks (e.g. channel levees). They were affected by episodic events such as floods but also by sedimentary infilling and channel mobility that played out over longer time scales.

The main fluvial harbour of Pisa (Tuscany, Italy) has been intensely investigated in recent decades, highlighting the importance of fluvial activity on the evolution of harbour structures (e.g. Benvenuti et al., 2006; Allinne et al., 2016). Pisa was an important Etruscan city

integrated into the Roman Republic during the 3rd century BC. The fluvial harbour “Stazione Pisa-San Rossore”, located close to the ancient city shows the constraints on fluvial harbours and the challenges that societies faced to adapt to these (Fig. 7A). According to Benvenuti et al. (2006) and Bruni (2000), the succession of floods led to the successive migration of the river course (palaeo-meander of the Serchio River) towards the Northwest and led to the destruction of numerous installations and ships. This channel was used as a harbour from the 5th century BC until the 5th or the 6th centuries AD, before its complete




Fig. 7. (A) Geomorphological map of the Pisan deltaic plain and location of the different harbours (adapted from Bini et al., 2015; Allinne et al., 2016; Kaniewski et al., 2018). B. Regional location of the site. C. Location of the palaeo-channels of the Serchio and the Arno rivers around the Etruscan/Roman city of Pisa (Prehistoric channels are denoted in yellow, Etruscan/Roman channels in light blue and Mediaeval channels in light gray; adapted from Bini et al., 2015; Image Microsoft Bing Maps). D–E. Landscape reconstruction showing two evolutionary stages of the lagoon of Portus Pisanus as documented by historical sources and archaeological data (adapted from Kaniewski et al., 2018). (D) Roman period – a wide lagoon basin, hosting Portus Pisanus as mentioned in literary sources. E. Late Middle Ages – the accretion of arcuate beach ridges, belonging to the Arno Delta strandplain, led to the closure of the lagoon. Construction of the maritime harbour of Livorno in a seaward position with respect to the lagoon. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

infilling and eventual abandonment (Camilli and Setari, 2005). Several flood-damaged wrecks have been dated between the 2nd century AD and the 5–6th centuries AD. It seems clear that the harbour basins were relocated in tandem with the migration of the channel towards the north-west. This translates the adaptation of the inhabitants to river migration more than to flood risk (Allinne et al., 2016). The city of Pisa was strongly affected by river floods due to the dense and complex network of fluvial channels related to the Arno and Serchio rivers. Bini et al. (2015) have identified three different river channels in the city area (Fig. 7.C) and have shown that, in Etruscan and Roman times (5th century BC – 2nd century AD), inhabitants expanded Pisa's urban network following the spatial distribution of topographically elevated areas, mainly formed by a complex and unstable fluvial network.

In a similar vein, the wreck of a Roman barge discovered in the Rhone River at Arles (Bouches-du-Rhone, France) further attests to the importance of floods on ancient fluvial harbours. Vella (2011) and Vella et al. (2014) have described erosion surfaces typical of an increase in fluvial dynamics at the top of cores drilled below the wreck. This erosion surface is followed by a coarse sedimentation that culminated with the sinking of the boat around 50–66 AD (Vella, 2011). The wreck, lying between 4 and 9 m depth, was overlain by fluvial sediments, including ceramics from the 1st/2nd centuries AD (Long, 2009a). A

longitudinal cross-section undertaken in the Grand Rhone, at the wreck site, underlines an artificial bank elevation of around 3.2 m during the 1st century AD (Mellinand and Sivan, 2011). The bank was reinforced at this time, possibly due to increasing harbour activity in this area of the river (Vella et al., 2014). The stability of the bank is attested by the 19th century docks that are located above the Roman bank (Vella et al., 2009).

Geoarchaeological investigations by Giaime et al. (2019) reveal that the settlement of Halmyris (Danube Delta, Romania) exploited a secondary palaeo-channel of the Saint-George arm of the Danube River during Antiquity (Fig. 8). The presence of this channel, characterized by chironomids and ostracods living in flowing waters, partially explains the location of the Getic settlement (6th – 1st centuries BC), followed by the Roman fortress (1st – 7th centuries AD). Due to its sheltered position, Halmyris was protected from Danube flooding. The disconnection of the Halmyris channel from the main Saint-George branch of the Danube appears contemporaneous with the abandonment of the settlement during the 7th century AD.

These examples present different aspects of flood hazard management during Antiquity. Pisa's harbour engineers chose to build light structures, which could be easily rebuilt when the channel shifted course. Archaeological excavations undertaken in the fluvial basins




Fig. 8. (A). Geomorphological map of the Danube delta close to Halmyris. The figure highlights the favourable location of Halmyris, on an elevated area and close to the palaeo-channel where the harbour was located in Getic (6th – 1st centuries BC) and Roman (1st– 7th centuries AD) times (after [Giaime et al., 2019](#)). B. Regional location of the site.

have demonstrated that the artificial structures were composed of wood and no “hard” harbour installations (quays, artificial banks) have been discovered ([Allinne et al., 2016](#)). At Arles, however, a flood management strategy seems to have been preferred. This scheme is reflected by the stabilisation of the riverbank during the 1st century AD. Around 2000 years ago, the hydrological activity of the Rhone was important. [Arnaud-Fassetta and Landuré \(2014\)](#) have demonstrated that the mean discharge was high and several high-energy flood episodes have been recorded. Nevertheless, the number of overbank flooding events affecting the city was reduced after the elevation of the banks ([Bruneton et al., 2001](#); [Vella et al., 2014](#)). By contrast, Halmyris had a slightly elevated position compared to the surrounding delta plain and was naturally protected from Danube floods. Furthermore, in comparison to Arles and Pisa, the harbour was built in a low-energy river channel. The main pressure on the harbour of Halmyris was subsequently linked to the siltation and the migration of the channel.

3.2. River-mouth harbours

Fluvial harbours are/were used by vessels adapted to fluvial navigation, which transported goods that were often unloaded in river-mouth harbours ([Boetto, 2010](#)). Since Early Antiquity, fluvial mouths have been used as safe havens, notably during storms. The use of river mouths as shelters dates back to the Bronze Age at least, with possible examples on the coast of Israel including Nahariya, Tel Achziv, Tel Akko or Tel Abu Hawam ([Raban, 1985](#)). However, over the long term, such environments were vulnerable to a plethora of natural hazards, both of marine and terrestrial origin ([Anthony, 2009](#)). Hydrological, sedimentological, biological and human processes interact and can induce domino effects on the environment. The infilling and mobility of a

river-mouth environment can have an upstream or a downstream origin because such environments are affected by fluvial discharge and marine-sediment deposition, even in the Mediterranean's micro-tidal regime. The vulnerability of river-mouth harbours to numerous hazard types explains why the cluster analysis did not clearly isolate an estuarine group within the dataset. River-mouth harbours are clearly delineated in Group A (Tel Akko – Bronze and Iron Age) which demonstrates the impact of sedimentary inputs into lagoonal-estuary environments, where this anchorage was located. We also identified two river-mouth harbours in the “fluvial group” (Group C) related to the importance of fluvial activity in their management (Castérou/Mandirac and Ostia). The importance of RSL changes in the geomorphological evolution of two river-mouth harbours is also discernible for the submerged sites of Herakleion and Les-Saintes-Maries-de-la-Mer found in Group B.

[Raban's \(1985\)](#) hypothesis for Levantine harbours is confirmed for the harbours of Tel Akko (Haifa Bay, Israel). Imported archaeological material (Egypt, Cyprus, the Aegean) demonstrates that the city must have had a harbour since the Bronze Age ([Artzy and Beeri, 2010](#); [Artzy, 2012](#); [Artzy et al., submitted](#)). At that time, a lagoon-type estuary was used as an anchorage area ([Morhange et al., 2016a](#); [Giaime et al., 2018](#)). Bio-sedimentological analyses have shown that this environment was permanently connected to the sea and constituted a good natural anchorage area. The gradual closure of the lagoon, probably driven by the morphogenesis of a sand spit and its infilling by sediments transported by the Na'aman River, engendered the relocation of the anchorage on the open coast (the western façade of the tell in Late Persian/Early Hellenistic times).

Geoarchaeological research undertaken in Ostia (mouth of the Tiber River, Italy) reveals the close relationships between the city and the




Fig. 9. A. Palaeogeography of Ostia's harbour system in Antiquity (adapted from Goiran et al., 2010, 2014; Vittori et al., 2015; Salomon et al., 2017, 2018). The Ostia river-mouth basin was abandoned before the construction of Portus in the 1st century AD. The figure highlights the importance of the Tiber delta's progradational dynamics since Roman times; B. Regional location of the site.

river since its foundation during the late 4th/early 3rd centuries BC (Salomon et al., 2017, 2018; Fig. 9). Significant deltaic progradation associated with fluvial changes since Roman times led to the progressive infilling of the channel and of the related harbour facilities. Located in the immediate proximity of the castrum in the 4th century BC, the channel migrated to the north in the 1st century BC (Salomon et al., 2017). Although two small harbour basins have been identified along the river at Ostia, it seems that most of the city's riverbanks were used as a harbour canal (Goiran et al., 2014; Sadori et al., 2016; Salomon et al., 2018). In one of the fluvial channels, an abrupt facies change between the pre-harbour facies and the harbour facies, associated with a chronological gap, suggests that the harbour basin was possibly excavated 6 m below Roman sea level and adapted for large maritime ships (Goiran et al., 2014). Nonetheless, heavy tonnage boats were unable to navigate up the river to the harbour because of offshore bars at the mouth of the Tiber (Boetto, 2010; Goiran et al., 2014; Salomon et al., 2014). As early as the 1st century AD, the basin was infilled by flood deposits (Goiran et al., 2014; Sadori et al., 2016) and the water column was reduced to 0.5 m rendering the harbour obsolete (Goiran et al., 2014). The construction of Portus' harbour complex (Claudius' basin) in the central part of the delta, inaugurated by the Emperor Nero in 64 AD, overcame the environmental challenges of the river-mouth harbour at Ostia. This new deltaic harbour allowed the area to retain its role as a leading port city in Roman Imperial times.

The importance of rivers, and their associated lagoons, has been highlighted for the Roman city of Narbonne (Aude, France; Sanchez et al., 2014, Sanchez et al., submitted; Faisse et al., 2018). The ancient Bubresus lake (nowadays Bages-Sigean lagoon; Fig. 10) constituted an ideal natural environment to accommodate a harbour. The lagoon was protected from the sea and had a water depth of at least 3 m in Early Roman times (Faisse et al., 2018; Sanchez et al., submitted). This

explains the high density of Roman settlements located on the palaeo-islands and the waterfront of the lagoon. In the Augustan period (27 BC – 14 AD), 50 sites (of the 124 archaeological sites identified for the period) were (30 sites) or could have been (20 sites) used for harbour purposes (Carayon et al., 2017). The fluvial harbour of Narbonne could be accessed by boat through the southern arm of the Aude River (that roughly corresponds to the present position of the Robine Canal). The first major installations on the channel mouth are dated to the 1st century AD (Sanchez et al., submitted). Two parallel artificial jetties, ~2 km long, were built on fluvial sandbanks (Faisse et al., 2018; Fig. 10 C–D). The two jetties (15–25 m wide) were ~50 m apart and built to maintain a 3.5–4 m deep channel (Mathé et al., 2016; Faisse et al., 2018; Sanchez et al., submitted). They were equipped with quays to unload and transfer goods from large maritime vessels to shallow-draught fluvial ships, or to be transported by terrestrial roads. Because of the intense fluvial sedimentary inputs, Roman engineers reduced the width of the northern part of the channel in order to increase the flow velocity and avoid siltation of the channelized course (Faisse et al., 2018; Sanchez et al., submitted). Archaeological excavations along these jetties have demonstrated that they were probably extended several times in order to keep the entrance of the channel clear of sediment. The fragility of these structures is attested by numerous repairs during the Early Roman Empire and explains the construction of a breakwater on the eastern side of the channel, to protect the area from swell (Sanchez et al., submitted).

Group B incorporates two river-mouth harbours (Herakleion, once located in the former Canopic mouth of the Nile and Les-Saintes-Maries-de-la-Mer, today submerged off the coast of the Rhone Delta). These harbours are characterized by high fluvial sediment supply that led to the formation of deltaic lobes where the harbours were hosted. RSL changes and the silting of the former channels led to the eventual




Fig. 10. A. Geomorphology of the Bages-Sigean lagoon complex. The red square denotes the location of the Castérou/Mandirac river-mouth facilities (adapted from Faïsse et al., 2018). B. Regional location of the site; C-E. Palaeogeographical reconstruction of the Castérou/Mandirac area before the 1st century AD (C), in the 1st century AD (D) when the first structures were built, and in the 5th century AD (E) at the time of the abandonment of the site (adapted from Faïsse et al., 2018). (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)




Fig. 11. A. Palaeogeography of the Saint-Feréol channel of the Rhone Delta (France) in Roman times (adapted from Pagès et al., 2008; survey Luc Long). B. Regional location of the site. C. Scenario for the progradation of the Rhone delta during the past 5000 years (adapted from Vella et al., 2005).

destruction of the lobes.

Geomorphological investigations on the Rhone Delta (France) have shown that, in Antiquity, the delta's coastline was locally more advanced in the sea, due to the location of the mouth of the Saint-Feréol arm of the river (Vella et al., 2005; Arnaud-Fassetta and Provansal, 2014). Seaward, the discovery of several ancient wrecks allowed researchers to reconstruct the morphology of the palaeo-mouth of the ancient Rhone of Saint-Feréol (Fig. 11). The wrecks were grounded on off-shore bars formed at the mouth of the river whilst trying to enter the channel (Long, 2009b). The Saint-Feréol branch of the Rhone was one of the major arms of the river in Antiquity and numerous ancient deltaic settlements lay on the banks of the channel (Arnaud-Fassetta and Landuré, 2014). One important anchorage seems to have been located in this part of the delta between the 5th century BC and the 5th century AD. Long and Duperron (2011) have excavated anchors and limestone blocks, up to 10 m below present mean sea level, interpreted as possible relics of harbour constructions (Long, 2009b). The subsidence of these harbour structures is associated with an erosion of the deltaic lobe induced by a strong reduction in sediment supply following the upstream

infilling of the Saint-Feréol branch that started during the second part of the 1st century AD (Arnaud-Fassetta and Landuré, 2014).

Shoreline retreat and deltaic subsidence are important in explaining the evolution of the Canopic mouth of the Nile River (Western Nile Delta, Egypt), where the ancient harbours cities of Herakleion and Eastern-Canopus were located (Fig. 12). Archaeological and geophysical investigations have highlighted the presence of harbour structures and shipwrecks up to 6 m below present mean sea level (Goddio, 2010; Hamouda et al., 2015; Robinson and Goddio, 2015). “Sediment failure” has been evoked to explain this subsidence, including the cumulative impacts of earthquakes, Nile floods or high-energy events, in addition to human constructions on unconsolidated sediments (Stanley et al., 2004, 2006). By contrast, recent work by Flaux et al. (2017a) has elucidated two periods of negative sediment budget at the Canopic mouth. The first, between 5500 and 4000 BP, was linked to a decrease in Nile discharge. The second phase occurred during the last 500 years, leading to a significant decrease in the sedimentary budget of the Canopic lobe and an erosion of the promontory. Relative sea-level rise linked to the compaction and liquefaction of unconsolidated muds culminated in the




Fig. 12. A. Geomorphological features of the Abu Qir Bay/Canopic coast on the western Nile Delta (adapted from Flaux et al., 2017a). The bathymetry of the western bay and the positions of Herakleion and East Canopus are adapted from Goddio (2007). B. Regional location of the site.

complete submersion of the promontory. The morphology of the Canopic river mouth, a promontory extending out into the sea, made it particularly exposed to high-energy events. Flaux et al. (2017a) identified two marine intrusions, recorded in the Canopic lobe's closed-lagoon environments, dated to the 1st and 2nd-3rd centuries AD that could correspond to high-energy events having affected Alexandria (Goiran, 2012).

The different examples highlighted above show that river-mouth harbours are located in challenging geomorphological contexts. In the short term, both fluvial floods and marine storms can affect them. At longer timescales, fluvial sediment budgets play a major role in shaping their geomorphological evolution (Flaux et al., 2017a). The two submerged estuarine harbours, Herakleion and Les-Saintes-Maries-de-la-Mer, highlight that these environments are extremely sensitive to rapid

relative sea-level rise and a decrease in sediment supply. Nevertheless, the vast majority of ancient river-mouth harbours are today landlocked on deltaic plains. Good examples include the harbours of Ravenna and Classe (Emilia Romagna, Italy; Cirelli, 2013; Manzelli, 2000) separated from the sea by the morphogenesis of a large coastal spit, or the harbour of Lattara (Languedoc-Roussillon, France) whose decline was partially induced by the intra-lagoonal progradation of the Mosson and Lez Rivers (Sabatier et al., 2010).

3.3. Lagoonal harbours

Lake and lagoon environments were particularly attractive locations for harbours during Antiquity (see Morhange et al., 2015, 2017 for further references). On the basis of our statistical analyses, most of the




Fig. 13. A. The ancient Bosphorus of the Taman Peninsula, south-east Russia (adapted from Giaime et al., 2016). B. Regional location of the site.

lagoonal harbours are found in group A (Cumes-Licola, Pollentia, Orgame, Portus Pisanus, the inner harbour of Seleucia Pieria). Included in this group is the unique example of Maryut lagoon's ancient harbours (e.g. Portus Mareoticus, Taposiris Magna, Marea-Philoxenia). Cumulatively, these port systems constituted the lacustrine harbour of Alexandria, and were located in a mixed lagoonal system that was affected by seasonal Nile floods (Group C). We demonstrate that these environments, while being naturally protected from the sea, were relatively short-lived due to long-term problems of harbour accessibility.

Lagoons, isolated from the sea by coastal spits formed by longshore drift processes, are an important component of Holocene deltas. Such a landscape is still visible on the Danube Delta (Romania), namely the large Razelm-Sinoe lagoon complex, or at a larger scale in the Kuban Delta (Taman peninsula, Russia). Today, lagoons constitute around 6400 km² of the Mediterranean coastline (Cataudella et al., 2015), but their surface areas were even more significant in Antiquity. In fact, due to base-level sedimentary inputs, most have been totally or partially infilled (Boyd et al., 1992; Oertel et al., 1992; Nichols and Boon, 1994; Duck and Figueiredo da Silva, 2012). This phenomenon is particularly evident on the Kuban Delta where various authors have reconstructed the transformation of the former Taman archipelago into a peninsula, shaped by the progressive infilling of the former Bosphorus channels linking the Black and the Azov Seas (Brückner et al., 2010; Kelterbaum et al., 2011; Fouache et al., 2012; Giaime et al., 2014 and Giaime et al., 2016; Fig. 13).

Founded in the 7th century BC, the ancient city of Orgame is located on the shores of the Razelm-Sinoe lagoon (southern Danube Delta, Romania). This important lagoon was formed by the reworking of Danube sediments, transported in particular by the Saint-George and Dunavăț arms of the river and reworked by longshore drift to form several generations of coastal spits that progressively isolated the lagoon from the sea (Giosan et al., 2006; Vespremeanu-Stroe et al., 2017; Fig. 14). The isolation of the lagoon has been elucidated using sedimentary cores extracted from the harbour basin of Orgame (Bony et al., 2015). The presence of a natural inlet, allowed the populations to

exploit a calm water body directly connected to the sea. The large surface area of the lagoon explains why the anchorage area of Orgame was not rapidly infilled. The transformation of the area into a phragmites wetland, and the infilling of the harbour, only began around 1000 years ago because of the influence of the Dunavăț pro-delta (Bony et al., 2015). The ancient harbour of Orgame is therefore unique in the history of lagoonal harbours in the Mediterranean/Black Sea world because most of the lagoonal harbours have been infilled and land-locked since Antiquity.

Kaniewski et al. (2018) have elucidated that the harbour of Portus Pisanus (the main harbour of Pisa) located north of the city of Livorno (Tuscany, Italy) was hosted in a marine lagoon that gradually evolved into a coastal lake (Fig. 7.D–E). The development of the protected lagoon started around 200 ± 30 BC in relation to the formation of a sand spit, fed by fluvial sediment reworked by the longshore currents. In Roman times, the lagoon water body was well protected and had a good connection with the sea, which made the area perfect for maritime transport and trade. There is no evidence of environmental changes until 1000 ± 20–1250 ± 20 AD when the lagoon started to lose its connection to the sea because of coastal progradation. The port was completely isolated from the sea in the late 13th century AD.

Numerous ancient lagoonal harbours were rapidly infilled by sediments during Antiquity, leading to the loss of harbour functions for many settlements. This explains why lagoon harbours in deltaic contexts were generally used for a short period of time. The formation of the lagoon of Cumes-Licola (Campania, Italy) resulted from the growth of a sand spit built by the sediments of the Volturno River, to the north of the Cumes palaeo-promontory (Fig. 15). This lagoon constituted a good environment for a harbour because it was naturally protected from the sea. However, geomorphological investigations have demonstrated that, since Hellenistic times, the lagoon shows signs of rapid infilling because of the erosion of surrounding hillslopes linked to the development of agriculture (Stefaniuk et al., 2003; Stefaniuk and Morhange, 2010). Further infilling led to an important reduction in the water depth of the lagoon. With a depth of < 1 m from Late Antiquity




Fig. 14. A-B. Palaeogeographical reconstruction of Orgame's coastal landscapes ~3500–2000 yr cal. BP (A) and ~1200 yr cal. BP (B) (adapted from Bony et al., 2015). C. Regional location of the site. The erosion of the promontory to the north of Orgame is attested by the destruction of ancient structures located on the cliff (ca. 20 m in height).



Fig. 15. A. Geomorphology of the southern margin of the Volturno Delta and location of the ancient harbour of Cumes in the infilled lagoon of Licola. High sediment supply is attested by the large sand spit which has isolated the lagoon from the sea (adapted from Morhange et al., 2015). B. Regional location of the site.

onwards, the lagoon was no longer accessible to ships.

The ancient lagoon of *Pollentia* (Mallorca, Balearic Archipelago, Spain), believed to have hosted one of the city's ancient harbours, underwent a similar evolution (Giaime et al., 2017; Fig. 16). The palaeo-

lagoon is located downstream of the northern littoral cell of Alcúdia Bay, north of the mouth of the Muro fluvial system. Due to its shallow water depth, the lagoon was probably dredged at the time of its foundation in the 1st century BC. Nevertheless, its accessibility remained limited and maritime ships would have been anchored offshore, near the Alcanada islet, some 4 km from the lagoon. The use of lighters, between the outer harbour and the lagoon harbour, linked the two areas.

Lake Maryut (Mareotis), which lies behind Alexandria on the western margin of the Nile Delta, hosted a great diversity of harbour structures during Antiquity (Fig. 17). Archaeological surveys undertaken on the shores of the Maryut have demonstrated that a large part of the lagoon was equipped with jetties related to artisanal or agricultural sites and that several main harbours were located on the lake's waterfront (Blue et al., 2011; Rodziewicz, 2011). At Taposiris Magna (Fig. 17.B), access to the Roman harbour was restricted by funnel-shaped entrance and exit areas, linked by a 1.7-km-long channel (Boussac and El-Amouri, 2010; Tronchère et al., 2012). The harbour of Marea-Philoxenia comprises a 2-km-long pier intersected by several quays constructed at right angles to the shoreline to form separate basins. These docking areas were active between the 5th and 7th centuries AD, for pilgrimage and trading purposes. The considerable length of the quays made year-round docking possible despite seasonal differences in the Maryut's water level (up to 1.5 m) driven by Nile floods. The fluvio-lagoonal system of the Maryut was divided into different geographical and trading areas, with the lacustrine harbour of Alexandria (Portus Mareoticus) being the focal point. Recent work by Flaux et al. (2017b) has highlighted the archaeological potential of Alexandria's lacustrine waterfront in Roman times. Combining bio-sedimentological analyses of sedimentary archives, the study of ancient maps and historical descriptions of the area (mainly from Strabo; Strabo 17.7), Flaux et al. (2017b) have highlighted the presence of two areas probably related to Portus Mareoticus (Fig. 18). They identified at least three ancient jetties, perpendicular (nowadays buried below the present urban fabric of Alexandria) to the lake's shoreline. These structures are several hundred meters long and were used from the 1st century BC to Late Roman times.

These examples demonstrate that the size of the lagoon and the importance of sedimentary input by local rivers are key factors in shaping the continuity of the harbours installed in such contexts. The reconstructed history of Portus Pisanus shows that societies successfully exploited the natural potentiality of a coastal lagoon to host an important harbour system from the 2nd century BC to the 11-12th centuries AD. The lagoons of Cuma-Licola and Pollentia highlight the role of limiting factors (i.e. limited accommodation space) that can affect the accessibility and the navigability of smaller lagoons. At Pollentia, the lagoon was perfectly located, at the foot of the city, but its depth, despite dredging, never allowed large maritime ships to moor there. For the Cuma-Licola lagoon, the mobility of the inlet, associated with the width of the sand spit and the presence of nearshore bars suggests that it was not ideal for naval circulation and for the installation of sustainable harbour activity throughout Antiquity. This explains why many of the region's harbours were located far from the Volturno River, such as Pozzuoli, Baiiae or Misenum (Morhange et al., 2015). Maryut lagoon constitutes an interesting case of harbour duality, with both fluvial and lagoonal characteristics. It necessitated specific facilities to accommodate intra-lagoonal navigation. The lagoon decreased in size after its isolation from the Nile and the Mediterranean Sea, from the 9th century AD onwards (Flaux et al., 2012). Settlement of the Mareotis waterfront decreased considerably from the 8th century AD onwards. Nonetheless, no environmental causal link has been directly established and this abandonment could be related to the progressive decline of the use of fluvial transport in addition to a decrease in the political importance of Alexandria after the Arab conquest.




Fig. 16. A. Geomorphological map of the ancient lagoon of Pollentia, Mallorca (adapted from Gaiame et al., 2017). The harbours were located in the northern part of the lagoon. The ancient inlet was probably located close to the present inlet. B. Regional location of the site.

3.4. Artificialisation of the harbours

Numerous Roman harbours have been densely artificialized which demonstrates the importance of human activities in maintaining harbour access routes and links to the open sea. During the Roman period, this often necessitated considerable port infrastructure. One example is the dense human artificialization of the Aude river-mouth to stabilize its riverbanks for port activities (Fig. 10). The Group G determined by the cluster analysis also includes densely equipped harbour basins. These harbours are characterized by the importance of human constructions and the dredging operations needed to maintain a sufficient water depth for maritime circulation in a context of shoreline progradation. In addition, we detailed the case of Alexandria's maritime harbours that were densely artificialized during Late Hellenistic/Roman times but that are nowadays submerged due to the subsidence of the Nile Delta front (Group B). In Magnus Portus (Eastern Harbour), in particular, a plethora of ancient port structures are now submerged. During the Hellenistic-Roman period, harbour infrastructure was diversified. An artificial diversion of one of the arms of the Küçük Meanders River was ordered by Hadrian in the Early 2nd century AD, in order to prevent the siltation of the harbour channel (Kraft et al., 2011; Stock et al., 2016).

During the 1st-2nd centuries BC, the discovery of hydraulic mortar led to a revolution in the construction of harbours along clastic coasts

(Brandon et al., 2014). Hydraulic mortar was probably discovered in the Campi Flegrei region of Italy, and was used for the harbours of the Naples Gulf, and then later for *Portus*, the harbour complex of Rome during the 1st century AD (Fig. 9). Initiated by Claudius, the construction of the first basin started in 42 AD in order to provide Rome with a maritime harbour, directly linked to the city via the Tiber River. Fifty years later, Trajan furnished the harbour with a second basin dug to the east of the Claudian basin. The Claudian basin, due to its location in the centre of the Tiber Delta front, was directly (i) exposed to wave action, (ii) subjected to river floods and (iii) to rapid sedimentation. The *Portus* complex was connected to the Tiber via several channels (Salomon et al., 2014, 2016; Lisé-Pronovost et al., 2018). The depth of the Claudius basin was around 6–8 m at the time of its construction and it was adapted to the mooring of large maritime ships (Goiran et al., 2010, 2011). Sedimentary cores have revealed that the sediments trapped in the harbour were mostly sandy, testifying to a large opening to the sea and high-energy dynamics. Despite the impressive moles that protected the harbour basin (Fig. 9), this facies differs from the common harbour facies found in protected harbours (Marriner and Morhange, 2006a). This hypothesis is reinforced by the writings of Tacitus who describes the destruction of around 200 ships following a storm in 62 AD (Tacitus, *Annals*, 15.18.2: “There was no addition to the price, although about two hundred ships were destroyed in the harbour by a violent storm, and one hundred more, which had sailed up the Tiber,




Fig. 17. A. Hydro-geomorphological map of the north-west Nile Delta during Antiquity, showing the location of sites discussed in the text (adapted from Flaux et al., 2017b). B. Regional location of the site. C. Artificial constructions at Taposiris harbour (adapted from Boussac and El-Amouri, 2010; Image: Microsoft Bing Maps).

by an accidental fire”). This situation could explain Trajan’s decision to build a second inner basin (Trajan’s harbour) to provide better anchorage for boats (Goiran et al., 2010). Goiran et al. (2011) have revealed, by coring the channel connecting the two basins, a reduction in the energy towards Trajan’s basin, marked by a decrease in the sediment grain size. The basin resembles a lagoon environment directly linked to the river and to the sea. The abandonment of Trajan’s harbour is documented by an infilling of the channel between the end of the 2nd century AD and the 5th century AD whereas, at that time, the Claudian basin was still navigable with a 5-m water column (Goiran et al., 2011). The example of Portus perfectly illustrates the importance of hydraulic mortar in facilitating the construction of large waterfront harbours. It explains the significant number of harbours built using this method since the 2nd century BC in the Mediterranean (Brandon et al., 2014).

The artificial structures of the ancient harbours of Alexandria also highlight the importance of Pozzolana. After Alexander the Great’s

conquest of Egypt and the foundation of Alexandria in 331 BC, two harbour basins were constructed on the leeward side of Pharos Island (Fig. 18). This area was perfect for maritime harbours because it was partially protected by the sandstone ridge (Pharos Island) that acted as a natural breakwater. The existence of the two harbours is attested on either side of the tombolo formed after the construction of an artificial causeway (*Heptastadium*) linking Pharos Island to the continent (Goiran, 2005; Marriner et al., 2008). The Eastern harbour (Magnus Portus) is known to have been densely artificialized during the Roman period, as recounted by Strabo (*Geography*, 17.1.6–17) who explains that the Eastern Bay was divided into several harbour basins (Hohlfelder and Oleson, 2014). The constructions continued after Strabo’s visit of Alexandria, as confirmed by the structures in pumiceous ash from Pozzolana and dated to the 1st century AD, currently drowned in the basin (Hohlfelder and Oleson, 2014). Harbour structures are nowadays submerged in the Eastern basin (Goddio and Yoyotte, 2008;






Fig. 18. A. Geomorphological map of Alexandria (Western Nile Delta, Egypt) and the location of its ancient harbours. The figure highlights the accretion of the tombolo after the construction of the artificial causeway (Heptastadion) following the foundation of the city by Alexander the Great in 331 BC (adapted from Goiran, 2001; Marriner et al., 2008; Flaux et al., 2017b). B. Regional location of the site.

Goddio and Masson-Berghoff, 2016). Stanley et al. (2006) have suggested that the submersion of these structures was probably accentuated by their construction on unconsolidated coastal sediments. The cumulative impact of natural hazards, such as earthquakes, storm

waves and tsunamis probably also played a role in the destabilisation of the sediments.

In deltaic contexts, Pozzolana structures have been used to protect harbours. The harbour of *Forum Iulli*, in the modern city of Frejus (Var,



- Estérel and Les Maures massif
- Holocene fluvial deposits
- Dune
- Beaches
- Modern lagoon
- Roman harbour

(caption on next page)

Fig. 19. A. Geomorphological map of the Argens Delta and location of the harbour basin of Frejus on the distal margin of the 11-km-long Holocene ria (adapted from Dubar, 2004). The positions of the successive coastlines in the lower valley are adapted from Bertonecello et al., 2014 (black line: Neolithic coastline; black dashed line: Bronze Age coastline; black dotted line: first Iron Age coastline; red dotted line: second Iron Age coastline; red dashed line: coastline at 0BC/AD; red line: Early Roman Empire coastline). They identified the succession of several beach-ridges and lagoonal environments similar to the current coastal morphology since the Bronze Age. B. Map of the Roman city of Forum Iulii with the location of the structures including volcanic tuff and Pozzolana material (after Excoffon and Dubar, 2011). The position of the coastline in the 1st century AD is from Excoffon et al. (2006) and the position of the coastline in the 1st century AD is from Gébara and Morhange, 2011 (adapted from Bony et al., 2011). C. Regional location of the site. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

France) is a landlocked harbour presently located ca. 1.5 km from the coast. The harbour was first mentioned in 31 BC (Tacitus repeated by Pliny the Elder, *Annals* 4.5), when Frejus harbour hosted part of Antony and Cleopatra's fleet, captured by Octavian after the battle of Axium. No archaeological structures related to the harbour are earlier than the Augustan period (end of the 1st c. BC – beginning of the 1st c. AD; Gébara and Morhange, 2011). In Roman times, Frejus was located on the eastern limit of the Argens Ria, at the palaeo-mouth of the Argens and Reyran Rivers (Fig. 19). Due to its position, at the margin of the Argens-Reyran river mouth, the harbour experienced problems of silting (Excoffon et al., 2010) that necessitated regular dredging (Bony et al., 2011). In order to control the sedimentary infilling of the basin, inhabitants also built a jetty to prevent sediments from filling the harbour entrance. Excoffon and Dubar (2011), have highlighted the use of pozzolana in the construction of this jetty, built in the continuity of the southern mole of Frejus harbour (Fig. 19).

4. The geoarchaeological trajectory of ancient deltaic harbours

The geomorphological trajectories of ancient harbours in deltaic contexts show that early coastal societies favoured natural anchorages along the coast (lagoons, river-mouths). Independently of any deltaic influence, such naturally protected areas are scarce, which necessitated the development of early harbour engineering. Marriner et al. (2014) have reported the early artificialisation, beginning in the Bronze Age, of pocket beaches along the coast of the Levant. The examples of Tyre, Sidon and Beirut clearly show the evolution of harbour infrastructure in their sediment records (Ancient Harbour Parasequence; Marriner and Morhange, 2006b; Marriner et al., 2014). In deltaic contexts, the utilisation of naturally protected areas did not require such man-made harbour protection systems, at least in Early Antiquity. Furthermore, rapid deltaic progradation since the Bronze Age did not allow the perpetuation of harbour activities in such areas. This means that the different sediment facies are not vertically stacked because of the relocation of the harbour basins through time, to keep pace with the shifting delta front. At Ephesus (Western Anatolia), intensive geoarchaeological work undertaken during the past two decades reveals the succession of at least six different harbours from the Late Bronze to Late Byzantine periods (Fig. 20). The first anchorages (Sacred and Koresos harbours) had few man-made structures. This is attested by the absence of a classic harbour sequence (Marriner and Morhange, 2006b), the sediments comprising marine sands that are consistent with a progradational sequence (Kraft et al., 2007; Stock et al., 2014). Use of these areas – since the 11th century BC for the Sacred harbour and since the mid-8th century for the Koresian harbour – are attested by ceramics and the contemporaneous multiplication of riverine settlements along the coast, showing a clear parallel between harbour location and settlement patterns. The same link is visible at Tel Akko (Haifa Bay, Israel; Artzy, 2012; Morhange et al., 2016a; Giaime et al., 2018; Artzy et al., submitted). In this maritime trade city, largely occupied (and urbanized) from the Middle Bronze Age onwards, the settlement pattern is linked to fluvio-coastal changes and harbour displacement. Former harbour activities were concentrated in the lagoonal estuary of the Na'aman River

during the Bronze and Iron Ages before their relocation to the western façade during Persian times. Subsequently, sometime between the late 3rd or the beginning of the 2nd century BC, the tell was abandoned following the relocation of the harbour on the Akko promontory, 1.5 km away and beyond the influence of fluvial processes. The absence of artificial structures at the two sites is explained by the natural protection of the anchorages.

At Ephesus, the harbours' location at the back of the transgressed ria meant that they were well protected from coastal storms. At Akko, the lagoonal-estuary was well sheltered. At both Akko and Ephesus, the artificial harbour constructions are attested from the Hellenistic period onwards. At Akko, a mole was constructed to protect the harbour basin from the swell after the relocation of the harbour in the Akko promontory, far from the fluvial influence (Giaime et al., 2018). The harbour was then extended in Roman and Medieval times, because Akko was an important harbour city and the Hellenistic basin is still in use today (Galili et al., 2010; Galili and Rosen, 2017). The heart of the Anatolian city was definitively moved and intensive constructions started on the western side of Mount Pion during the Hellenistic period. At the same time, the first harbour structures seem to have been constructed. The artificial mole built during the middle-Hellenistic period was lengthened and possibly reinforced at several stages during Roman times (Kraft et al., 2007). Nonetheless, the widespread human interventions (constructions and dredging) on the harbour basin were not sufficient to prevent the siltation of the basin, whose accessibility was drastically reduced in Late Roman and Byzantine times. In the end, the harbour could only be accessed by small, shallow-draught vessels. Maritime boats had to be unloaded in natural anchorages downstream.

Deltaic contexts did not significantly affect the evolution of harbour structures. By contrast, harbours and their associated settlements had to regularly be relocated to keep abreast with deltaic progradation, which has led to the term “race to the sea” being coined.

5. Conclusion

This typology has shed light on the evolution of ancient harbour basins in deltaic contexts. Each geomorphological context presents specific advantages and disadvantages. The earliest harbours were hosted in naturally protected areas. In deltaic contexts, river mouths were preferred, but such environments were poor long-term harbours because of the combined action of sedimentary inputs, river changes and high-energy fluvial and marine events. Nonetheless, ancient populations developed techniques to improve or maintain the natural potentialities of their anchorage areas, before the discovery of hydraulic mortar and the construction of more grandiose harbour complexes along coastlines lacking natural anchorages.

Despite floods, storms and sedimentary inputs, human populations adapted to these changing and, at times, challenging environments. Nevertheless, human responses to environmental modifications were highly variable. The “race to the sea” seems to have been the norm in Antiquity, sometimes leading to the complete abandonment of settlements and their relocation in areas less vulnerable to deltaic progradation.




Fig. 20. A. Progradation of the Küçük Menderes (Cayster) deltaic plain since the Neolithic and B. Regional location of the site. C. Location of the successive anchorages of Ephesus, the Sacred Harbour (1), Koressos harbour (2), the Roman and Byzantine harbour (3), the Byzantine harbour (4), the Late Byzantine and later harbour (5), Pygela harbour (6) (adapted from Brückner, 2005; Stock et al., 2013).

Acknowledgements

M. Giaime acknowledges the support of the Institute of Advanced Studies and the Department of Geography at Durham University and was supported by a Durham Junior Research Fellowship, cofunded by the European Union under grant agreement number 609412 (MSCA COFUND). The project leading to this publication received funding from the Excellence Initiative of Aix-Marseille Université–A*MIDEX, a French ‘Investissements d’Avenir’ project in the framework of Labex OT-Med (ANR-11-LABEX-0061).

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.earscirev.2019.01.022>.

References

Tacitus, Annals 13–16, with English translation by J. Jackson (1937), Loeb Classical Library, London and Cambridge, Mass.
 Tacitus, Annals 4, with English translation by J. Jackson (1937), Loeb Classical Library, London and Cambridge, Mass.
 Strabo, Geography of Strabo. Jones, H.L., 1969, Geography of Strabo in Eight Volume (Vol. VIII). Harvard University Press, Cambridge, Massachusetts (517 pp.).

Secondary literature

Allinne, C., 2007. Les villes romaines face aux inondations. La place des données archéologiques dans l'étude des risques fluviaux. *Géomorphol. Relief Process. Environ.* 13 (1), 67–84.
 Allinne, C., Morhange, C., Pasquinucci, M., Roumieux, C., 2016. Géoarchéologie des ports de Pise « Stazione Ferroviaria San Rossore » et de *Portus Pisanus*: dynamiques géomorphologiques, sources antiques, données archéologiques. *Revue Archéologique de Narbonnaise* 44, 321–338 Supplément.
 Amorosi, A., Bini, M., Giacomelli, S., Pappalardo, M., Ribecai, C., Rossi, V., Sammartino, I., Sarti, G., 2013. Middle to late Holocene environmental evolution of the Pisa coastal plain (Tuscany, Italy) and early human settlements. *Quat. Int.* 303, 93–106.
 Anthony, E.J., 2009. *Shore Processes and their Palaeoenvironmental Applications*. Elsevier.
 Anthony, E.J., Marriner, N., Morhange, C., 2014. Human influence and the changing geomorphology of Mediterranean deltas and coasts over the last 6000 years: from progradation to destruction phase? *Earth Sci. Rev.* 139, 336–361.
 Arnaud-Fassetta, G., Landuré, C., 2014. Le risque fluvial en milieu rural de l'époque grecque au haut Moyen Âge, Le cas du delta du Rhône (France), in N. Carcaud, G. Arnaud-Fassetta (dir.), *La géoarchéologie française au XXI^{ème} siècle*, 215–236. CNRS Editions, Paris.
 Arnaud-Fassetta, G., Provansal, M., 2014. The lower valley and the delta of the Rhône River: water landscapes of nature and history. In: Fort, M., André, M.M. (Eds.), *Landscapes and Landforms of France*, pp. 207–218 (Dordrecht).
 Arnaud-Fassetta, G., Carre, M.B., Marocco, R., Maselli-Scotti, F., Pugliese, N., Zaccaria, C., Bandelli, A., Bresson, V., Manzoni, G., Montenegro, M.E., Morhange, C., Pipan, M., Prizzon, A., Siché, I., 2003. The site of Aquileia (northeastern Italy): example of fluvial geoarchaeology in a Mediterranean coastal plain. *Géomorphol. Relief Process. Environ.* 9 (4), 227–245.
 Arnaud-Fassetta, G., Carcaud, N., Castanet, C., Salvador, P.G., 2010. Fluvial palaeoenvironments in archaeological context: geographical position, methodological approach and global change-hydrological risk issues. *Quat. Int.* 216, 93–117.
 Artzy, M., 2012. Return to Tel Akko, Its Anchorages, Harbor, and Surroundings. *Leon Recanati Institute for Maritime Studies Newsletter*, pp. 5–14.
 Artzy, M., Beeri, R., 2010. Tel Akko. In: Killebrew, A.E., Raz-Romero, V. (Eds.), *One Thousand Nights and Days – Akko Through the Ages*. Hecht Museum, Haifa, pp. 14–23.
 Artzy M., Morhange C., Giaime M., Jol H., Lopez G.I., Abu-Hamid A., Salmon Y. and Killebrew A.K., Coastline, river changes and their effects on anchorages / harbors and habitation patterns: Akko (Israel) as an example, In: Blue L., Sivan D., Farr H., Baika K. and Blackman D., (Eds.), *Harbours and Maritime Cultural Landscapes*, 2019 submitted, Sidestone Press; Leiden.
 Bellotti, P., Caputo, C., Davoli, L., Evangelista, S., Garzanti, E., Pugliese, F., Valeri, P., 2004. Morpho-sedimentary characteristics and Holocene evolution of the emergent part of the Ombrone River delta (southern Tuscany). *Geomorphology* 61, 71–90.
 Bellotti, P., Calderoni, G., Carboni, M.G., Di Bella, L., Tortora, P., Valeri, P., Zernitskaya, V., 2007. Late Quaternary landscape evolution of the Tiber River delta plain (Central Italy): new evidence from pollen data, biostratigraphy and 14C dating. *Z. Geomorphol.* 51 (4), 505–534.
 Benvenuti, M., Mariotti-Lippi, M., Pallechhi, P., Sagri, M., 2006. Late-Holocene catastrophic floods in the terminal Arno River (Pisa, Central Italy) from the story of a Roman riverine harbour. *The Holocene* 16 (6), 863–876.
 Bertonecello, F., Devillers, B., Bonnet, S., Guillon, S., Boubry, L., Delhon, C., 2014. Mobilité des paysages littoraux et peuplement dans la basse vallée de l'Argens (Var, France) au

cours de l'Holocène. *Quaternaire* 25 (1), 43–44.
 Bini, M., Rossi, V., Amorosi, A., Pappalardo, M., Sarti, G., Noti, V., Capitani, M., Fabiani, F., Gualandi, M.L., 2015. Palaeoenvironments and palaeotopography of a multi-layered city during the Etruscan and Roman periods: early interaction of fluvial processes and urban growth at Pisa (Tuscany, Italy). *J. Archaeol. Sci.* 59, 197–210.
 Blackman, D.J., 1982a. Ancient harbours in the Mediterranean, part 1. *Int. J. Naut. Archaeol.* 11, 79–104.
 Blackman, D.J., 1982b. Ancient harbours in the Mediterranean, part 2. *Int. J. Naut. Archaeol.* 11, 185–211.
 Blue, L., Khalil, E., Trakadas, A. (Eds.), 2011. *A Multidisciplinary Approach to Alexandria's Economic Past: the Lake Mareotis Research Project*. British Archaeological Report. International Series, 2285 & Series in Archaeology Monographs (University of Southampton) Vol. 5 Archaeopress, Oxford.
 Boetto, G., 2010. Le port vu de la mer: l'apport de l'archéologie navale à l'étude des ports antiques. *Bollettino Archeol. Online B/B7/9*, 112–128. Online. Available from: http://www.bollettinodiarcheologiaonline.beniculturali.it/documenti/generale/9_Boetto.paper.pdf.
 Bony, G., Morhange, C., Bruneton, H., Carbonel, P., Gébara, C., 2011. 2000 ans de colmatage du port antique de Fréjus (Forum Julii), France: une double métamorphose littorale. *Compt. Rendus Geosci.* 343 (10), 701–715.
 Bony, G., Morhange, C., Marriner, N., Baralis, A., Kaniowski, D., Rossignol, I., Lungu, V., 2015. History and influence of the Danube delta lobes on the evolution of the ancient harbour of Orgame (Dobrogea, Romania). *J. Archaeol. Sci.* 61, 186–203.
 Boussac, M.F., El-Amouri, M., 2010. The lake structures at Taposiris. In: Blue, L., Khalil, E. (Eds.), *Lake Mareotis. Reconstructing the Past*. British Archaeological Reports. International Series, 2113 & Series in Archaeology Monographs (University of Southampton), 2 Archaeopress, Oxford, pp. 87–105.
 Boyd, R., Dalrymple, R., Zaitlin, B.A., 1992. Classification of clastic coastal depositional environments. *Sediment. Geol.* 80, 139–150.
 Brandon, C.J., Hohltdfelder, R.L., Jackson, M.D., Oleson, J.P., 2014. *Building for Eternity. The History and Technology of Roman Concrete Engineering in the Sea*. Oxbow Books, Oxford.
 Brückner, H., 2005. Holocene shoreline displacements and their consequences for human societies: the example of Ephesus in Western Turkey. *Zeitschrift für Geomorphologie* 137, 11–22 Supplement-Volume.
 Brückner, H., Müllenhoff, M., Gehrels, R., Herda, A., Knipping, M., Vött, A.V., 2006. From archipelago to floodplain – geographical and ecological changes in Miletus and its environs during the past six millennia (Western Anatolia, Turkey). *Zeitschrift für Geomorphologie* 142, 63–83 Supplement-Volume.
 Brückner, H., Kelterbaum, D., Marunchak, O., Porotov, A., Vött, A., 2010. The Holocene Sea level story since 7500 BP — lessons from the Eastern Mediterranean, the Black and the Azov Sea. *Quat. Int.* 225 (2), 160–179.
 Brückner, H., Herda, A., Müllenhoff, M., Rabbel, W., Stümpel, H., 2014. On the Lion Harbour and other Harbours in Miletos: recent historical, archaeological, sedimentological, and geophysical research. In: *Rune Frederiksen Soren Handb (Ed.), Proc. Danish Inst. Athens*. Vol. 7. pp. 49–103.
 Brückner, H., Herda, A., Kerschner, M., Müllenhoff, M., Stock, F., 2017. Life cycle of estuarine islands—from the formation to the landlocking of former islands in the environs of Miletos and Ephesos in western Asia Minor (Turkey). *J. Archaeol. Sci. Rep.* 12, 876–894.
 Bruneton, H., Arnaud-Fassetta, G., Provansal, M., Sistach, D., 2001. Geomorphological evidence of fluvial change during the roman period in the lower Rhône valley. *Catena* 45, 287–312.
 Bruni, S., (dir.), 2000. *Le navi antiche di Pisa / The ancient ships of Pisa: ad un anno dall'inizio delle ricerche / after a year of work*. Polistampa, Firenze.
 Camilli, A., Setari, E., 2005. *Le navi antiche di Pisa*. Guida Archeologica. Electa, Milano (dir. 63).
 Camuffo, D., Secco, C., Brimblecombe, P., Martin-Vide, J., 2000. Sea storms in the Adriatic Sea and the western Mediterranean during the last millennium. *Climate Change* 46, 209–223.
 Carayon, N., Keay, S., Arnaud, P., Sanchez, C., 2017. The harbour system of Narbo Martius (Narbonne/F) and its facilities during antiquity. In: von Carnap-Bornheim, C., Daim, F., Ettl, P., Warnke, U. (Eds.), *Harbours as Objects of Interdisciplinary Research – Archaeology + History + Geosciences*, pp. 151–164 (Verlag des Römisch-Germanischen Zentralmuseums).
 Carsana, V., Febbraro, S., Giampaola, D., Guastaferro, C., Irollo, G., Ruello, M.R., 2009. Evoluzione del paesaggio costiero tra Parthenope e Neapolis. *Méditerranée* 112, 14–22.
 Casana, J., 2008. Mediterranean valleys revisited: linking soil erosion, land use and climate variability in the Northern Levant. *Geomorphology* 101 (3), 429–442.
 Cataudella, S., Crosetti, D., Massa, F. (Eds.), 2015. *Mediterranean coastal lagoons: sustainable management and interactions among aquaculture, capture fisheries and the environment*. FAO, Rome General Fisheries Commission for the Mediterranean, Studies and Reviews 95, Rapport de la.
 Catuneanu, O., Zecchin, M., 2013. High-resolution sequence stratigraphy of clastic shelves II: controls on sequence development. *Mar. Pet. Geol.* 39 (1), 26–38.
 Chelbi, F., Paskoff, R., Troussat, P., 1995. La baie d'Utique et son évolution depuis l'Antiquité: Une ré-évaluation géoarchéologique. *Antiq. Afr.* 31, 7–51.
 Gianflone, G., Cavuoto, G., Punzo, M., Dominici, R., Sonnino, M., Di Fiore, V., Pelosei, N., Tarallo, D., Lirer, F., Critelli, S., De Rosa, R., 2018. Late quaternary stratigraphic setting of the Sibari Plain (southern Italy): Hydrogeological implications. *Marine and Petroleum Geology* 97, 422–436.
 Cirelli, E., 2013. Roma sul mare e il porto augusteo di Classe. In: Boschi, F. (Ed.), *Ravenna e l'Adriatico dale origini all'età romana*. Ante Quem, Bologna, pp. 109–121.
 Dawson, A.G., Stewart, I., 2007. Tsunami deposits in the geological record. *Sediment. Geol.* 200, 166–183.

- Day Jr., J.W., Gunn, J.D., Folan, W.J., Yanez-Arancibia, A., Horton, B.P., 2007. Emergence of complex societies after sea level stabilized. *Eos Trans. AGU* 55 (15), 169–170.
- Dearing, J.A., Jones, R.T., 2003. Coupling temporal and spatial dimensions of global sediment flux through lake and marine sediment records. *Glob. Planet. Chang.* 39 (1–2), 147–168.
- Delile, H., Abichou, A., Gadhoun, A., Goiran, J.P., Pleuger, E., Monchambert, J.Y., Wilson, A., Fentress, E., Quinn, J., Ben Jerbania, I., Ghazzi, F., 2015. The geoarchaeology of Utica, Tunisia: the paleogeography of the Mejerda Delta and hypotheses concerning the location of the ancient harbor. *Geoarchaeology* 30 (4), 291–306.
- Devillers, B., 2008. Holocene Morphogenesis and Anthropisation of a Semi-Arid Watershed, Gialias river, Cyprus. BAR International, Series 1775 Archaeopress, Oxford.
- Di Donato, V., Ruello, M.R., Liuzza, V., Carsana, V., Giampaola, D., Di Vito, M.A., Morhange, C., Cinque, A., Russo Ermolli, E., 2018. Development and decline of the ancient harbor of Neapolis. *Geoarchaeology* 33, 542–557.
- Dubar, M., 2004. L'édification de la plaine deltaïque du Bas Argens (Var, France) durant la Protohistoire et l'Antiquité. Application d'un modèle numérique 2D à l'archéologie. *Méditerranée* 102 (1), 47–54.
- Duck, R.W., Figueiredo da Silva, J., 2012. Coastal lagoons and their evolution: a hydro-morphological perspective. *Estuar. Coast. Shelf Sci.* 110, 2–14.
- Dusar, B., Verstraeten, G., Notebaert, B., Bakker, J., 2011. Holocene environmental change and its impact on sediment dynamics in the Eastern Mediterranean. *Earth Sci. Rev.* 108 (3–4), 137–157.
- Engel, M., Brückner, H., 2011. The identification of palaeo-tsunami deposits – a major challenge in coastal sedimentary research. *Coastline Rep.* 17, 65–80.
- Engel, M., May, S.M., 2012. Bonaire's boulder fields revisited: evidence for Holocene tsunami impact on the Leeward Antilles. *Quat. Sci. Rev.* 54, 126–141.
- Excoffon, P., Dubar, M., 2011. Note sur l'emploi de tuf volcanique et de la pouzzolane dans quelques constructions de Forum Iulii. Eléments de réflexion sur l'utilisation et la diffusion de la pouzzolane en Méditerranée occidentale. *Revue du Centre Archéologique du Var* 2011, 171–181.
- Excoffon, P., Devillers, B., Bonnet, S., Bouby, L., 2006. Nouvelles données sur la position du littoral antique de Fréjus. Le diagnostic archéologique du « théâtre d'agglomération » (Fréjus, Var). *Archéosciences* 30, 205–221.
- Excoffon, P., Bonnet, S., Devillers, B., Berger, J.F., 2010. L'évolution du trait de côte aux abords de Fréjus, de sa fondation jusqu'à la fin du 1er siècle après J.-C. *Archéologie des rivages méditerranéen* 50, 47–54.
- Faïsse, C., Mathé, V., Bruniaux, G., Labussière, J., Caverio, J., Jézégou, M.P., Lefèvre, D., Sanchez, C., 2018. Palaeoenvironmental and archaeological records for the reconstruction of the ancient landscape of the Roman harbour of Narbonne (Aude, France). *Quat. Int.* 463, 124–139.
- Flaux, C., Marriner, N., El-Assal, M., Morhange, C., Rouchy, J.M., Soulié-Marsche, I., Torab, M., 2012. Environmental changes in the Maryut lagoon (northwestern Nile delta) during the last ~2,000 years. *J. Archaeol. Sci.* 39 (12), 3493–3504.
- Flaux, C., Marriner, N., El-Assal, M., Kaniewski, D., Morhange, C., 2017a. Late Holocene erosion of the Canopic promontory (Nile Delta, Egypt). *Mar. Geol.* 385, 56–67.
- Flaux, C., El-Assal, M., Shaalan, C., Marriner, N., Morhange, C., Torab, M., Goiran, J.P., Empereur, J.Y., 2017b. Geoarchaeology of Portus Mareoticus: Ancient Alexandria's lake harbour (Nile Delta, Egypt). *J. Archaeol. Sci. Rep.* 13, 669–681.
- Fouache, E., Kelterbaum, D., Brückner, H., Lericolais, G., Porotov, A., Dikarev, V., 2012. The late Holocene evolution of the Black Sea — a critical view on the so-called Phanagorian regression. *Quat. Int.* 266 (2012), 162–174.
- Gallili, E., Rosen, B., 2017. The Akko Marina Archaeological Project— Summary. In: Gallili, E. (Ed.), *The Akko Marina Archaeological Project*. BAR International Series 2862 BAR Publishing, Oxford, England, pp. 320–344.
- Gallili, E., Rosen, B., Zviely, D., Silberstein, N., Finkielsteyn, G., 2010. The evolution of Akko harbour and its Mediterranean maritime trade links. *J. Island Coast. Archaeol.* 5, 191–211.
- Gébara, C., Morhange, C., 2011. Fréjus (Forum Iulii): le port antique/the ancient harbour. *J. Roman Archaeol.* 77 supplement series. Oxford.
- Giaime, M., Baralis, A., Marriner, N., Morhange, C., Porotov, A., 2014. D'un archipel à une plaine deltaïque: géoarchéologie de la Péninsule de Taman (Sud-Ouest de la Russie). *Géochronique* 130, 42–45.
- Giaime, M., Avnaim-Katav, S., Morhange, C., Marriner, N., Rostek, F., Porotov, A.V., Baralis, A., Kaniewski, D., Brückner, H., Kelterbaum, D., 2016. Evolution of Taman Peninsula's ancient Bosphoruses, south-West Russia: deltaic progradation and Greek colonization. *J. Archaeol. Sci. Rep.* 5, 327–335.
- Giaime, M., Morhange, C., Ontiveros, M.Á.C., Fornós, J.J., Vacchi, M., Marriner, N., 2017. In search of Pollentia's southern harbour: geoarchaeological evidence from the Bay of Alcúdia (Mallorca, Spain). *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 466, 184–201.
- Giaime, M., Morhange, C., Marriner, N., López-Cadavid, G.I., Artzy, M., 2018. Geoarchaeological investigations at Akko, Israel: new insights into landscape changes and related anchorage locations since the Bronze Age. *Geoarchaeology* 33 (6), 641–660.
- Giaime, M., Magne, G., Bivolaru, A., Gandouin, E., Marriner, N., Morhange, C., 2019. Halmyris: geoarchaeology of a fluvial harbour on the Danube Delta (Dobrogea, Romania). *The Holocene* 29 (2), 313–327.
- Giampaola, D., Carsana, V., Boetto, G., 2004. Il mare torna a bagnare Neapolis. Parte II: dalla scoperta del porto al recupero dei relitti. *L'archeologo subacqueo* 10, 15–19.
- Giosan, L., Donnelly, J.P., Constantinescu, S., Filip, F., Ovejanu, I., Vespremeanu-Stroe, A., Vespremeanu, E., Duller, G.A., 2006. Young Danube delta documents stable Black Sea level since the middle Holocene: morphodynamic, paleogeographic, and archaeological implications. *Geology* 34 (9), 757–760.
- Goddio, F., 2007. Topography and excavation of Heracleion-Thonis and East Canopus (1996–2006). Oxford Centre for Maritime Archaeology 1, Oxford (137 pp.).
- Goddio, F., 2010. Geophysical survey in the submerged Canopic region. In: Robinson, D., Wilson, A. (Eds.), *Alexandria and the North-Western Nile Delta*. Joint Conference Proceedings of Alexandria: City and Harbour (Oxford 2004) and Trade, Topography and Material Culture of Egypt's North-Western Delta (Berlin 2006). Oxford Centre for Maritime Archaeology Monograph 5, Oxford, pp. 3–13.
- Goddio, F., Masson-Berghoff, A. (Eds.), 2016. *The BP Exhibition: Sunken Cities: Egypt's Lost Worlds*. Thames and Hudson, London.
- Goddio, F., Yoyotte, J., 2008. The Portus Magnus of Alexandria. In: Goddio, F. (Ed.), *Egypt's Sunken Treasures*, 2nd ed. Prestel, London.
- Goff, J., Chagué-Goff, C., Nichol, S., Jaffe, B., Dominey-Howes, D., 2012. Progress in palaeotsunami research. *Sediment. Geol.* 243 (244), 70–88.
- Goiran, J.P., 2001. *Recherches géomorphologiques dans la région littorale d'Alexandrie en Égypte*. Thèse de doctorat en géographie physique, Université de Provence, Aix-Marseille I, Aix-en-Provence.
- Goiran, J.P., 2012. Caractérisation d'un dépôt de tsunami dans le port antique d'Alexandrie par l'étude exoscopique des quartz: apports et limites de la méthode. In: Rébé-Marichal, I., Laurenti, A., Boehm, I., Goiran, J.P. (Eds.), *Archéosismicité & Tsunamis en Méditerranée, Approches croisées*. Editions APS, Goulien, pp. 157–190.
- Goiran, J.P., Salomon, F., Tronchère, H., Djerbi, H., Carbonel, P., Ognard, C., Oberlin, C., 2001. Géoarchéologie des ports de Claude et de Trajan, Portus, delta du Tibre. *Mélanges de l'École française de Rome-Antiquité* 123 (1), 157–236.
- Goiran, J.P., Marriner, N., Morhange, C., el-Maguib, M.A., Espic, K., Bourcier, M., Carbonel, P., 2005. Geomorphological evolution of Alexandria's maritime façade during the past 6000 years. *Méditerranée* 104, 61–64.
- Goiran, J.P., Tronchère, H., Salomon, F., Carbonel, P., Djerbi, H., Ognard, C., 2010. Palaeoenvironmental reconstruction of the ancient harbors of Rome: Claudius and Trajan's marine harbors on the Tiber delta. *Quat. Int.* 216, 3–13.
- Goiran, J.P., Salomon, F., Mazzini, I., Bravard, J.P., Pleuger, E., Vittori, C., Boetto, G., Christiansen, J., Arnaud, P., Pellegrino, A., Pepe, C., Sadori, L., 2014. Geoarchaeology confirms location of the ancient harbour basin of Ostia (Italy). *J. Archaeol. Sci.* 41, 389–398.
- Hamouda, A., El-Gendy, N., El-Gharabawy, S., Salah, M., 2015. Acoustic survey along Herakleion and East Canopus ancient Greek cities, Abu Quir Bay, Alexandria, Egypt. *J. Earth Sci. Clim. Chang.* 6 (289). <https://doi.org/10.4172/2157-7617.1000289>. Available from:.
- Hohlfelder, R.L., Oleson, J.P., 2014. Roman maritime concrete technology in its Mediterranean context. In: Brandon, C.J., Hohlfelder, R.L., Jackson, M.D., Oleson, J.P. (Eds.), *Building for Eternity: The History and Technology of Roman Concrete Engineering in the Sea*. Oxbow Books, Oxford, pp. 223–235.
- Hori, K., Tanabe, S., Saito, Y., Haruyama, S., Nguyen, V., 2004. Kitamura A. Delta initiation and Holocene sea-level change: example from the Song Hong (Red River) delta, Vietnam. *Sedimentary Geology* 164 (3–4), 237–249.
- Kaniewski, D., Marriner, N., Morhange, C., Faivre, S., Otto, T., Van Campo, E., 2016. Solar pacing of storm surges, coastal flooding and agricultural losses in the Central Mediterranean. *Sci. Rep.* 6, 25197.
- Kaniewski, D., Marriner, N., Morhange, C., Vacchi, M., Sarti, G., Rossi, V., Bini, M., Pasquinnucci, M., Allinne, C., Otto, T., Luce, F., Van Campo, E., 2018. Holocene evolution of Portus Pisanus, the lost harbour of Pisa. *Sci. Rep.* 8, 11625.
- Portus: an archaeological survey of the Portus of imperial Rome. In: Keay, S., Millet, M., Parolli, L., Strut, K. (Eds.), *Archaeological Monographs of the British School at Rome*. Vol. 15 The British School at Rome, London.
- Kelterbaum, D., Brückner, H., Porotov, A., 2011. Geoarchaeology of Taman Peninsula (Kerch Strait, South-West Russia) — the example of the ancient Greek settlement of Golubitskaya 2. *Contrib. Geoarchaeol.* 3 (3), 235–258 DIE ERDE.
- Kennett, D., Kennett, J., 2006. Early state formation in southern Mesopotamia: Sea levels, Shorelines, and climate change. *J. Island Coast. Archaeol.* 1, 67–99.
- Kolaiti, E., Papadopoulos, G.A., Morhange, C., Vacchi, M., Triantafyllou, I., Mourtzas, N.D., 2017. Palaeoenvironmental evolution of the ancient harbor of Lechaion (Corinth Gulf, Greece): were changes driven by human impacts and gradual coastal processes or catastrophic tsunamis? *Mar. Geol.* 392, 105–121.
- Kraft, J.C., Bückner, H., Kayan, I., Engelmann, H., 2007. The geographies of ancient Ephesus and the Artemision in Anatolia. *Geoarchaeology* 22 (1), 121–149.
- Kraft, J.C., Rapp, G., Brückner, H., Kayan, I., 2011. Results of the struggle at ancient Ephesus: natural processes I, human intervention 0. *Geological Society London Special Publications* 35, 27–36.
- Lambeck, K., Antonioli, F., Purcell, A., Silenzi, S., 2004. Sea-level change along the Italian coast for the past 10,000 yr. *Quaternary Science Reviews* 23 (14–15), 1567–1598.
- Lambeck, K., Rouby, H., Purcell, A., Sun, Y., Sambridge, M., 2014. 2014. Sea level and global ice volumes from the Last Glacial Maximum to the Holocene. *Proceedings of the National Academy of Sciences* 111 (43), 15296–15303.
- Leveau, P., 2012. Attraits et risques de l'eau dans les sociétés antiques. Retour critique sur la problématique du risque fluvial. In J.P. Bost (dir.), *L'eau: usages, risques et représentations dans le Sud-Ouest de la Gaule et le Nord de la péninsule Ibérique, de la fin de l'Âge du Fer à l'Antiquité tardive (IIème av. J.-C. – VIème ap. J.-C.)*. *Aquitania*, Supplément 21, 26–48, Bordeaux.
- Lisé-Pronovost, A., Salomon, F., Goiran, J.P., St-Onge, G., Herries, A.L.R., Montero-Serrano, J.C., Heslop, D., Roberts, A.P., Levchenko, V., Zawadzki, A., Heijns, H., 2018. Dredging and canal gate technologies in Portus, the ancient harbour of Rome, reconstructed from event stratigraphy and multi-proxy sediment analysis. *Quat. Int.* <https://doi.org/10.1016/j.quaint.2018.05.018>.
- Long, L., Duperron, G., 2011. Sur les traces de l'avant-port maritime d'Arles. In: SFEACG, Actes du Congrès d'Arles, pp. 97–118 Arles.
- Long, L., 2009a. De la mer au fleuve les ports antiques d'Arles. In: Long, L., Picard, P. (Eds.), *César le Rhône pour mémoire*. Actes Sud, Musée départemental Arles Antique, Arles, pp. 30–43.
- Long, L., 2009b. *Bouches-du-Rhône. Au large des Saintes-Maries-de-la-Mer*. Bilan

- Scientifique – Direction Régionale des Affaires Culturelles Provence-Alpes-Côte d'Azur, Service Régional de l'Archéologie. Vol. 2009. pp. 54–55.
- Manzelli, V., 2000. Città romane. 2, Ravenna. Atlante Tematico di Topografia Antica. Supplementi, 7. L'Erma di Bretschneider, Roma.
- Marchetti, M., 2002. Environmental changes in the Central Po Plain (northern Italy) due to fluvial modifications and anthropogenic activities. *Geomorphology* 44, 361–373.
- Marriner, N., Morhange, C., 2006a. Geoarchaeological evidence for dredging in Tyre's ancient harbour. *Levant*. Quat. Res. 65, 164–171.
- Marriner, N., Morhange, C., 2006b. The 'Ancient Harbour Parasequence': anthropogenic forcing of the stratigraphic highstand record. *Sediment. Geol.* 186, 13–17.
- Marriner, N., Goiran, J.P., Morhange, C., 2008. Alexander the Great's tomolos at Tyre and Alexandria, Eastern Mediterranean. *Geomorphology* 100 (3–4), 377–400.
- Marriner, N., Flaux, C., Morhange, C., Kaniewski, D., 2012. Nile Delta's sinking past: quantifiable links with Holocene compaction and climate-driven changes in sediment supply? *Geology* 40 (12), 1083–1086.
- Marriner, N., Morhange, C., Kaniewski, D., Carayon, N., 2014. Ancient harbour infrastructure in the Levant: tracking the birth and rise of new forms of anthropogenic pressure. *Sci. Rep.* 4, 5554.
- Marriner, N., Kaniewski, D., Morhange, C., Flaux, C., Giaime, M., Vacchi, M., Goff, J., 2017a. Tsunamis in the geological record: making waves with a cautionary tale from the Mediterranean. *Sci. Adv.* 3 (10).
- Marriner, N., Morhange, C., Flaux, C., Carayon, N., 2017b. Harbors and ports, ancient. In: Gilbert, A.S. (Ed.), *Encyclopedia of Geoarchaeology*. Springer, Dordrecht, pp. 382–403.
- Maselli, V., Trincardi, F., 2013. Man made deltas. *Sci. Rep.* 3, 1926.
- Mathé, V., Sanchez, C., Bruniaux, G., Camus, A., Cavero, J., Faisse, C., Jézégou, M.P., Labussière, J., Lévêque, F., 2016. Prospections géophysiques multi-méthodes de structures portuaires antiques à Narbonne (Aude, France). *ArchéoSciences* (1), 47–63.
- Mellinand, P., Sivan, O., 2011. Gare Maritime, diagnostic. INRAP.
- Morhange, C., Marriner, N., 2010. Mind the (stratigraphic) gap: Roman dredging in ancient Mediterranean harbours. *Bollettino di Archeologia online* 1 (2010), 23–32 [Online] B/B7/9.
- Morhange, C., Pirazzoli, P.A., Evelpidou, N., Marriner, N., 2012. Late Holocene tectonic uplift and the silting up of Lechaion, the western harbor of ancient Corinth, Greece. *Geoarchaeology* 27 (3), 278–283.
- Morhange, C., Marriner, N., Blot, M.L., Bony, G., Carayon, N., Carmona, P., Flaux, C., Giaime, M., Goiran, J.P., Kouka, M., Lena, A., Oueslati, A., Pasquinucci, M., Porotov, A.V., 2015. Dynamiques géomorphologiques et typologie géoarchéologique des ports antiques en contextes lagunaires. *Quaternaire* 26 (2), 117–139.
- Morhange, C., Giaime, M., Marriner, N., Abu Hamid, A., Bruneton, H., Honnorat, A., Kaniewski, D., Magnin, F., Wante, J., Zviely, D., Artzy, M., 2016a. Geoarchaeological evolution of Tel Akko's ancient harbour (Israel). *J. Archaeol. Sci. Rep.* 7, 71–81.
- Morhange, C., Marriner, N., Carayon, N., 2016b. The eco-history of ancient Mediterranean harbours. In: Bekker-Nielsen, T., Gertwagen, R. (Eds.), *The Inland Seas. Towards an Ecohistory of the Mediterranean and the Black Sea*. Steiner Franz Verlag, Stuttgart, pp. 85–106.
- Morhange, C., Marriner, N., Bony, G., Flaux, C., Giaime, M., Kouka, M., 2017. Geoarchaeology of ancient harbours in lagoonal contexts: an introduction. In: Franconi, T. (Ed.), *Fluvial Landscapes in the Roman World*, *Journal of Roman Archaeology*, supplementary series 104 Portsmouth, Rhode Island, pp. 97–110.
- Morton, R.A., Gelfenbaum, G., Jaffe, B.E., 2007. Physical criteria for distinguishing sandy tsunami and storm deposits with modern examples. *Sediment. Geol.* 200, 184–207.
- Nichols, M.N., Boon, J.D., 1994. Sediment transport processes in coastal lagoons. In: Kjerfve, B. (Ed.), *Coastal Lagoon Processes*. Elsevier, Amsterdam, pp. 157–219.
- Oertel, G.F., Kraft, J.C., Kearney, M.S., Woo, H.J., 1992. A rational theory for barrier-lagoon development. In: Fletcher, C.H., Wehmler, J.F. (Eds.), *Quaternary Coasts of the United States*. SEPM, pp. 77–87 Special Publication 48.
- Pagès, G., Long, L., Fluzin, P., Dillmann, P., 2008. Réseaux de production et standards de commercialisation du fer antique en Méditerranée: les demi-produits des épaves romaines des Saintes-Maries-de-la-Mer (Bouches-du-Rhône). *Revue archéologique de Narbonnaise* 41 (1), 261–283.
- Pagliarulo, R., 2006. Coastal changes and the environmental evolution of the archaeological site of Sybaris (southern Italy). *Geogr. Fis. Din.* 29 (2), 51–59.
- Pilarczyk, J.E., Dura, T., Horton, B.P., Engelhart, S.E., Kemp, A.C., Sawai, Y., 2014. Microfossils from coastal environments as indicators of paleo-earthquakes, tsunamis and storms. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 413, 144–157.
- Pranzini, E., 2001. Updrift river mouth migration on cusped deltas: two examples from the coast of Tuscany, Italy. *Geomorphology* 38, 125–132.
- Pranzini, E., 2007. Airborne LIDAR survey applied to the analysis of the historical evolution of the Arno River delta (Italy). *J. Coast. Res.* SI50, 400–409.
- Quézel, P., Médail, F., 2003. *Ecologie et biogéographie des forêts du bassin méditerranéen*. Vol. 572 Elsevier, Paris.
- Raban, A., 1985. The ancient harbours of Israel in Biblical times. In: Raban, A. (Ed.), *Harbour Archaeology: Proceedings of the First International Workshop on Ancient Mediterranean Harbours, Caesarea Maritima, 24–28 June 1983*, pp. 11–44 (Bar International Series, 257, Oxford).
- Robinson, A., Goddio, F. (Eds.), 2015. *Thonis-Heracleion in Context*. Oxford Centre for Maritime Archaeology Monographs, Oxford.
- Rodziewicz, M.D., 2011. Wine production and trade in Late Roman Alexandria. In: Décobert, C., Empereur, J.-Y., Picard, C. (Eds.), *Alexandrie Médiévale*. 4. Etudes Alexandrines, 24 Centre d'Etudes Alexandrines, Alexandria, pp. 39–56.
- Ruiz, J.M., Carmona, P., Cueva, A.P., 2014. Flood frequency and seasonality of the Júcar and Turia Mediterranean rivers (Spain) during the "Little Ice Age". *Méditerranée* 122, 121–130.
- Sabatier, P., Dezileau, L., Barbier, M., Raynal, O., Lofi, J., Briquieu, L., Condomines, M., Bouchette, F., Certain, R., Von Grafenstein, U., Jorda, C., Blan-Chemanche, P., 2010. Late-Holocene evolution of a coastal lagoon in Gulf of Lions (South of France). *Bulletin de la Société Géologique de France* 181 (1), 27–36.
- Sadori, L., Mazzini, I., Pepe, C., Goiran, J.P., Pleuger, E., Ruscito, V., Salomon, F., Vittori, C., 2016. Palynology and ostracodology at the Roman port of ancient Ostia (Rome, Italy). *The Holocene* 26 (9), 1502–1512.
- Salomon, F., Goiran, J.P., Bravard, J.P., Arnaud, P., Djerbi, H., Kay, S., Keay, S., 2014. A harbour-canal at Portus: a geoarchaeological approach to the *Canale Romano*: Tiber delta, Italy. *Water Hist.* 6 (1), 31–49.
- Salomon, F., Keay, S., Strutt, K., Goiran, J.P., Millett, M., Germoni, P., 2016. Connecting Portus with Ostia: preliminary results of a geoarchaeological study of the navigable canal on the Isola Sacra. *Revue archéologique de Narbonnaise* 44, 293–304.
- Salomon, F., Goiran, J.P., Pannuzi, S., Djerbi, H., Rosa, C., 2017. Long-term interactions between the Roman City of Ostia and its Paleomeander, Tiber Delta, Italy. *Geoarchaeology* 32 (2), 215–229.
- Salomon, F., Goiran, J.P., Noirrot, B., Pleuger, E., Buckowiecki, E., Mazzini, I., Carbonel, P., Gadhoum, A., Arnaud, P., Keay, S., Zampini, S., Kay, S., Raddi, M., Ghelli, A., Pellegrino, A., Morelli, C., Germoni, P., Zampini, S., 2018. Geoarchaeology of the Roman port-city of Ostia: Fluvio-coastal mobility, urban development and resilience. *Earth Sci. Rev.* 177, 265–283.
- Sanchez, C., Jezegou, M.P. (dir.), 2011. *Espaces littoraux et zones portuaires de Narbonne et sa région dans l'Antiquité*. Monographies d'archéologie méditerranéenne, 28. Édition de l'Association pour le développement de l'archéologie en Languedoc-Roussillon, Lattes.
- Sanchez, C., Faisse, C., Jézégou, M.P., Mathé, V., 2014. Le système portuaire de Narbonne antique: approche géoarchéologique. In: Mercuri, L., González Villaescusa, R., Bertonecello, F. (Eds.), *Implantations humaines en milieu littoral méditerranéen: facteurs d'installation et processus d'appropriation de l'espace, de la Préhistoire au Moyen Age*, 34e rencontres internationales d'archéologie et d'histoire d'Antibes. Editions APDCA, Antibes, pp. 125–136.
- Sanchez C., Jézégou M.P., Cavero J., Faisse C., Mathé V., Flaux C., Camus A. and Bruniaux G., *The Ancient Port of Narbonne (France): Five Centuries of Fluvio-Lagoonal Development*, 2019 submitted.
- Simeoni, U., Corbau, C., 2009. A review of the Delta Po evolution (Italy) related to climatic changes and human impacts. *Geomorphology* 107, 64–71.
- Sivan, D., Wdowski, S., Lambeck, K., Galili, E., Raban, A., 2001. Holocene sea-level changes along the Mediterranean coast of Israel, based on archaeological observations and numerical model. *Palaeogeography, Palaeoclimatology, Palaeoecology* 167 (1), 101–117.
- Stanley, J.D., 1997. Mediterranean deltas: subsidence as a major control of relative sea-level rise. In: Briand, F., Maldonado, A. (Eds.), *Transformations and Evolution of the Mediterranean Coastline*. Bulletin du Musée océanographique de Monaco, n° special 18 CIEM Science Serie 3, Monaco, pp. 35–62.
- Stanley, J.D., Bernasconi, M.P., 2009. Sybaris-Thuri-Copia trilogy: three delta coastal sites become land-locked. *Méditerranée* 112, 75–86.
- Stanley, J.D., Bernasconi, M.P., 2012. Buried and submerged Greek archaeological coastal structures and artifacts as gauges to measure late Holocene seafloor subsidence off Calabria, Italy. *Geoarchaeology* 27 (3), 189–205.
- Stanley, J.D., Warne, A.G., 1994. Worldwide initiation of Holocene Marine deltas by deceleration of sea-level rise. *Science* 265, 228–231.
- Stanley, J.D., Warne, A.G., 1997. Holocene sea-level change and early human utilization of deltas. *Geol. Soc. Am. Today* 7, 1–7.
- Stanley, J.D., Goddio, F., Jorstad, T.F., Schnepp, G., 2004. Submergence of ancient Greek cities off Egypt's Nile delta. *Geol. Soc. Am. Today* 14 (1), 4–10.
- Stanley, J.D., Jorstad, T.F., Goddio, F., 2006. Human impact on sediment mass movement and submergence of ancient sites in the two harbours of Alexandria, Egypt. *Nor. J. Geol.* 86, 337–350.
- Stefaniuk, L., Morhange, C., 2010. Evoluzione dei paesaggi littorali nella depression sud-ovest di Cuma da 4000 anni, il problema del porto antico. In: Alessio, A., Lombardo, M., Siciliano, A. (Eds.), *Atti del 40 convegno di studi sulla Magna Grecia, Taranto (2008)*. Istituto per la storia e l'archeologia della Magna Grecia, Napoli, pp. 305–322.
- Stefaniuk, L., Brun, J.P., Munzi, P., Morhange, C., 2003. L'evoluzione dell'ambiente nei Campi Flegrei e le sue implicazioni storiche: il caso di Cuma e le ricerche del Centre Jean Bérard nella laguna di Licola. In: Stazio, A., Ceccoli, C. (Eds.), *Ambiente e paesaggio nella Magna Grecia: atti del quarantaduesimo Convegno di studi sulla Magna Grecia, Taranto, 5–8 ottobre 2002*. Istituto per la Storia e l'Archeologia della Magna Grecia, Taranto, pp. 397–435.
- Stewart, I.S., Morhange, C., 2009. Coastal geomorphology and sea-level change. In: Woodward, J. (Ed.), *The Physical Geography of the Mediterranean*. Oxford University Press, Oxford, pp. 385–413.
- Stiros, S., Pirazzoli, P., Rothaus, R., Papageorgiou, S., Laborel, J., Arnold, M., 1996. On the date of construction of Lechaion, western harbor of ancient Corinth, Greece. *Geoarchaeology* 11 (3), 251–263.
- Stock, F., Pint, A., Horejs, B., Ladstätter, S., Brückner, H., 2013. In search of the harbours: New evidence of Late Roman and Byzantine harbours of Ephesus. *Quat. Int.* 312, 57–69.
- Stock, F., Kerschner, M., Kraft, J.C., Pint, A., Frenzel, P., Brückner, H., 2014. The palaeogeographies of Ephesus (Turkey), its harbours, and the Artemision – a geoarchaeological reconstruction for the timespan 1500–300 BC. *Z. Geomorphol.* 58 (Supplementary issue 2), 033–066.
- Stock, F., Knipping, M., Pint, A., Ladstätter, S., Delile, H., Heiss, A.G., Laermanns, H., Mitchell, P.D., Ployer, R., Steskal, M., Thanheiser, U., Urz, R., Wennrich, V., Brückner, H., 2016. Human impact on Holocene sediment dynamics in the Eastern Mediterranean – the example of the Roman harbour of Ephesus. *Earth Surf. Process. Landf.* 41 (7), 980–996.
- Syvitski, J.P., Kettner, A., 2011. Sediment flux and the Anthropocene. *Philos. Trans. R.*

- Soc. Lond. A: Math. Phys. Eng. Sci. 369, 957–975.
- Tallet, P., Marouard, G., 2014. The Harbor of Khufu on the Red Sea Coast at Wadi al-Jarf, Egypt. *Near East. Archaeol.* 77 (1), 4–14.
- Tallet, P., Marouard, G., 2017. The Harbor Facilities of King Khufu on the Red Sea Shore: the Wadi al-Jarf/tell Ras Budran System. *J. Am. Res. Center Egypt* 52, 135–177.
- Teatini, P., Tosi, L., Strozzi, T., 2011. Quantitative evidence that compaction of Holocene sediments drives the present land subsidence of the Po Delta, Italy. *J. Geophys. Res.* 116 (B08407), 1–10 [Online].
- Törnqvist, T.E., Wallace, D.J., Storms, J.E., Wallinga, J., Van Dam, R.L., Blaauw, M., Derksen, M.S., Klerks, C.J.W., Meijneken, C., Snijders, E.M.A., 2008. Mississippi Delta subsidence primarily caused by compaction of Holocene strata. *Nat. Geosci.* 1 (3), 173–176.
- Tronchère, H., Millet, B., Goiran, J.P., Carbonel, P., Djerbi, H., Vera, R., Torab, M., El-Assal, M., Callot, Y., 2012. Geoaerchaeological results from the harbor of Taposiris and implications concerning the construction of the harbor. In: *Ägypten und Levante* 22–23, pp. 383–394.
- Vacchi, M., Marriner, N., Morhange, C., Spada, G., Fontana, A., Rovere, A., 2016. Multiproxy assessment of Holocene relative sea-level changes in the western Mediterranean: sea-level variability and improvements in the definition of the isostatic signal. *Earth Sci. Rev.* 155, 172–197.
- Vacchi, M., Ghilardi, M., Melis, R.T., Spada, G., Giaime, M., Marriner, N., Lorscheid, T., Morhange, C., Burjachs, F., Rovere, A., 2018. New relative sea-level insights into the isostatic history of the Western Mediterranean. *Quat. Sci. Rev.* 201, 396–408.
- Vella, C., 2011. Géomorphologie fluviale et archéologie subaquatique: un mariage naturel. In: Djaoui, D., Greck, S., Marlier, S. (Eds.), *Arles-Rhône 3, Le naufrage d'un chaland antique dans le Rhône, enquête pluridisciplinaire*. Actes Sud, Arles, pp. 204–207.
- Vella, C., Fleury, T.J., Raccasi, G., Provansal, M., Sabatier, F., Bourcier, M., 2005. Evolution of the Rhône delta plain in the Holocene. *Mar. Geol.* 222 (223), 235–265.
- Vella, C., Hermitte, C., Parisot, J.C., 2009. La résistivité électrique et la géomorphologie au service de l'Archéologie. In: Long, L., Picard, P. (Eds.), *César le Rhône pour mémoire*. Actes Sud, Arles, pp. 204–207 Musée départemental Arles Antique.
- Vella, C., Tomatis, C., Sivan, O., 2014. Le contexte paléoenvironnemental. In S. Marlier (dir.), *Arles-Rhône 3, un chaland gallo-romain du Ier siècle ap. J.-C.*, 65–73. Paris, CNRS éditions.
- Vespremeanu-Stroe, A., Zainescu, F., Preoteasa, L., Tatui, F., Rotaru, S., Morhange, C., Stanica, A., Hanganu, J., Timar-Gabor, A., Cardan, I., Piotrowska, N., 2017. Holocene evolution of the Danube delta: an integral reconstruction and a revised chronology. *Mar. Geol.* 388, 38–61.
- Vittori, C., Mazzini, I., Salomon, F., Goiran, J.P., Pannuzi, S., Rosa, C., Pellegrino, A., 2015. Palaeoenvironmental evolution of the ancient lagoon of Ostia Antica (Tiber delta, Italy). *Journal of Archaeological Science* 54, 374–384.
- Vött, A., 2007. Silting up Oiniadai's harbours (Acheloo River delta, NW Greece). Geoaerchaeological implications of late Holocene landscape changes. *Géomorphol. Relief Process. Environ.* 13 (1), 19–36.
- Vött, A., Schriever, A., Hand, M., Brückner, H., 2007. Holocene palaeogeographies of the central Acheloo River delta (NW Greece) in the vicinity of the ancient seaport Oiniadai. *Geodin. Acta* 20 (2007), 241–256.
- Wainwright, J., Thornes, J.B., 2004. Environmental issues in the Mediterranean: processes and perspectives from the past and present. Routledge.
- Weiss, R., 2012. The mystery of boulders moved by tsunamis and storms. *Mar. Geol.* 295 (298), 28–33.