



Coastal Archaeology and Climate Change in the Middle East and North Africa Contextualizing Global Projections

Kieran Westley and Georgia Andreou

Coastal archaeological features at Tell Ruqeish, dating to the first millennium BCE in conjunction with human interventions on the beach (recreational infrastructure, quarrying, terracing, waste). Photograph from the GAZAMAP project.

The impact of climate change on archaeological heritage has received increasing attention, reflected in an ever-growing number of academic studies, dedicated funding calls, and inclusion in national and international policy-influencing efforts. The consensus is that archaeological sites, monuments, and artifacts face a clear threat from climate change. The nature of the threat and its impact varies depending on the type of heritage in question and the prevailing weather and climate where it is located. However, it is generally understood that climate change will either exacerbate existing processes of long-term decay and weathering, or introduce new processes where they previously did not operate. In the worst cases, this could lead to significant damage to, or the destruction and loss of, irreplaceable archaeological material. Consequently, there is an urgent need to identify and implement sustainable ways to protect or mitigate damage to archaeological sites and landscapes (Hambrecht and Rockman 2017; ICOMOS 2019; Sesana et al. 2021).

This article focuses on maritime cultural heritage, with emphasis on coastal archaeology. Maritime archaeological sites provide testament to the engagement of people with the sea, offering opportunities to explore historical themes such as seafaring, migration, colonization, and international interaction, many of which continue to resonate in the modern world. In this respect maritime cultural heritage provides a vital link between

regional archaeologies, particularly in the Middle East and North Africa (MENA) region. It also offers paleoenvironmental evidence (e.g., sea-level change) with potential to inform global studies on climate change (e.g., Yasur-Landau et al. 2021).

Lying at the intersection of land and sea, coastal archaeology is particularly vulnerable to climate change, largely because coastal flooding and erosion can have a truly detrimental, long-term impact on maritime cultural heritage. Episodic flooding and erosion may also intensify locally depending on changing storm and tropical cyclone patterns and can result in rapid destruction, or major damage, within a few hours (Wright 2016; Gregory et al. 2022). Both phenomena are projected to increase incrementally over the twenty-first century in response to climate-change-driven sea-level rise (Vousdoukas et al. 2018, 2020). Although a precise global estimate does not presently exist, national and regional studies suggest that at least thousands of sites are vulnerable over the coming century (Anderson et al. 2017; Brooks et al. 2020; Li et al. 2022).

Because flooding and erosion are also hazards for coastal cities, communities, and infrastructure, an extensive body of coastal management research has modeled their future impacts (Hinkel et al. 2021). The broader applicability of this research has reached archaeologists and historic environment managers to varying degrees. These range from simple application of future flood levels (Anderson et al. 2017; Li et al. 2022), to more complex studies seeking to quantify the effects of flooding and erosion (Vousdoukas et al. 2022; Reimann et al. 2018). The recent development of global-scale, open-access geospatial models of




Figure 1. Archaeological sites in the Gaza Strip. Map adapted from Andreou et al. 2022a.

twenty-first-century erosion and flooding (Vousdoukas et al. 2018, 2020) has encouraged these latter types of studies. Such studies are an essential component of the archaeological response to climate change (Hambrecht and Rockman 2017). In theory, an inventory of vulnerable sites and associated frequency and intensity of climate-change-related threats can enable appropriate prioritization of responses and resources. Prioritization is particularly relevant to coastal archaeology, where it is now accepted that not all sites can be saved (Dawson et al. 2021). This is primarily because of the scale of the problem: There are simply not enough resources to deal with every site that will be flooded or eroded. Additionally, physically protecting sites, with sea defenses for example, is unsustainable in the long-term given rising sea-levels. This strategy also runs counter to coastal management strategies that, for retreating shorelines, often advise “managed realignment” (removal of coastal defenses to allow creation of natural intertidal or marsh areas) or “no active intervention,” particularly where there is no/little infrastructure to protect or where the ecological costs of protection outweigh the benefits (Williams et al. 2018).

In practice, the effectiveness of these models depends heavily on the quality of their inputs, including data availability and data resolution, all of which affect the scale of analysis and the questions that these models can answer. Thus, although the contribution of global models to broadscale threat identification has been generally received positively in maritime archaeology (Andreou et al. 2022b), they are not without limitations. Recent studies have highlighted biases in global and regional models of climate change, particularly the relatively limited consideration of human adaptations that potentially slow down impacts and whose implementation may depend on financial resources (Hinkel et al. 2021). Moreover, the necessary simplification of coastal processes when modeling on a global scale has raised concern because it can result in inaccurate or lower confidence projections of future change (Cooper et al. 2020). For example, it has been projected that by 2100 large parts of the coast of the UAE and Bahrain will advance despite sea-level rise (Vousdoukas et al. 2020). This conclusion results from the fact that the historic pattern of shoreline change used to inform the model is based on satellite imagery from 1984 onward: a period of massive and unprecedented




Figure 2. Eroding archaeological features on the coast of Tell Ruqeish, Gaza Strip. Photograph from the GAZAMAP project.



Figure 3. Recreational infrastructure carved on the coast of Tell Ruqeish, Gaza Strip defining the location of the modern coastline. Photograph from the GAZAMAP project.

land reclamation that has artificially extended the coastline. Therefore, the drivers for coastal advance in this context are more likely socioeconomic rather than climate-change-related and it is uncertain as to whether this trend will continue up to 2100, as modeled. This is not to say that projections are purposeless or ambiguous. Large-scale models are essential in providing information on general trends and impacts, which they do well in terms of the direction of future change. This can help raise awareness and, ideally, stimulate social, political, and economic actions to tackle the climate crisis. Large-scale models also provide the



foundation for more detailed studies that can examine the local impact of these general changes. These issues are, of course, known to the modeling community, who for coastal areas recommend detailed studies at smaller scales that shed light on the compounding natural and anthropogenic processes that drive change (Vousdoukas et al. 2022).

This is particularly important in the coastal zone, where multiple processes originating from the atmosphere, land, and sea interact with human activity to produce change over multiple spatial and temporal scales. The effect of coastal flooding and erosion can be influenced, for instance, by human actions that increase or decrease sediment supply—such as coastal construction, harbor development, and upstream damming—as much as by sea-level rise resulting purely from global warming (IPCC 2022). Although there are indisputable links between climate change and the deterioration of maritime archaeological material (Gregory et al. 2022), there is a conceivable risk that these are examined without actively considering associated human actions that could aggravate flooding and erosion, or that may have driven changes that are inaccurately interpreted as a direct consequence of climate change.

Here we take a critical stance toward the projected impact of climate change on maritime archaeology with emphasis on coastal heritage. We argue that finding the root cause of coastal impacts such as flooding or coastal erosion can be difficult because both can be created by climate-change-related sea-level rise, unrelated human actions, or an overlapping combination

of the two. Additional modifiers include physical processes such as waves, tides, and storms that also exhibit natural variability but are themselves being modified by climate change. Nonetheless, identifying drivers of change is useful to understand the future trend of impacts and ultimately the most appropriate adaptation or mitigation response for a given location.

We illustrate this through case studies from the Gaza Strip, Libya, and Oman. These areas are currently subject to in-depth study by the authors within the framework of a MENA-wide maritime archaeological threat assessment: the Maritime Endangered Archaeology (MarEA) project (Andreou et al. 2022b). Moreover, these areas all have multiple data sources ranging from satellite imagery to local knowledge that suggest that a combination of physical processes and anthropogenic actions has impacted coastal archaeological sites. We then discuss the implications of this complexity in terms of projections of future impact, management considerations, and mitigation strategies. Finally, we highlight the contribution of archaeology to wider climate change research.

Coastal Erosion in the Gaza Strip

The southern Levant benefits from an impressive array of scientific studies exploring long-term environmental dynamics

and their intersection with maritime archaeology (Yasur-Landau et al. 2021). This reflects both the suitability of this region for such studies, but also underscores significant investment and expertise in the study and preservation of heritage that is not necessarily available along the broader Levantine coast. A case in point is the Gaza Strip, widely discussed in conjunction with interlinked ecological, humanitarian, and economic crises, all of which contribute to the rapid deterioration of cultural heritage.

As is the case at multiple Levantine coastal sites (e.g., Arsuf, Ashkelon, Byblos, and Yavne-Yam: Galili and Rosen 2010; Deroin, Bou Kheir, and Abdullah 2017; Barkai et al. 2018) coastal erosion is affecting Gaza's maritime archaeology. Sites on the coast of Rafah, near the Iron Age site of Tell Ruqeish, and around the multiperiod site of Anthedon have lost over one meter of land every year between the 1990s and 2021 (Klinger and Knauer 2020; Andreou et al. 2022a; figs. 1–2). Such rates are generally viewed as a twentieth-century phenomenon, and the tendency has been to consider erosion greater than one-half meter per year as anthropogenic, associated, among others, with the modernization of traditional land-use practices and the construction of dams, modern harbors, and offshore structures (Anthony, Marriner, and Morhange 2014). This interpretation does not exclude the possibility of premodern erosion, for which the evidentiary record is limited but potentially significant (e.g. Marriner et al. 2012). Rather, it highlights the lack of environmental studies accompanying modern alterations, the impact of which appears more extensive. In this respect, there is abundant and relatively easily accessible evidence for erosion, including imagery from the 1950s onward (Anthropocene), while evidence from the Holocene relies on geoarchaeological studies within the small-scale of archaeological sites.

Along the Levantine coast, rapid coastal retreat has been linked to decreased deposition of Nile-sourced sediment caused by construction of the Aswan Dam in the 1970s (Abd-el Monsef, Smith, and Darwish 2015). Unfortunately, the lack of reliable pre-1970s aerial imagery for Gaza (Andreou et al. 2022a) prevents a reconstruction of pre-Aswan coastal trends that would enable a systematic comparison of recent versus past erosion rates. At the same time, smaller-scale coastal assessments link erosion primarily with the construction of modern harbors and subsequently with agricultural activities, recreational development, and conflict-related damage (Abualtayef et al. 2012, 2021; Andreou et al. 2022a).

Erosion rates exceeding one meter per year are projected to continue, according to global models that rely on post-1980s satellite imagery (Vousdoukas




Figure 6. Comparison of recent (post-2010) and historic (pre-2010) shoreline movements at Apollonia generated from high resolution recent satellite images (Maxar, provided by the European Space Agency), declassified KH-9 images (provided by the USGS) and historic air photos. Colored lines show shoreline movement rates (LRR) in meters/year. Bottom image shows excerpt from the historic map of Beechey and Beechey (1828: 536).




Figure 7. Comparison of 1974 KH-9 (provided by the USGS) and 2016 satellite images (Maxar, provided by the European Space Agency) west of Ptolemais showing extensive retreat of waterline (i.e., beach loss) and backshore cliff erosion. Scars on the beach are suggestive of sand mining.

et al. 2020). Though it is reasonable to suggest that climate change has increased the rate of erosion in Gaza, it is not possible to view it independently from extensive human interventions on the coastal landscape (fig. 3). It is also not possible, at present, to quantify long-term land loss, owing to the


unknown impact of unpredictable human activities on the coast. Though landscape alterations are evident, the erosion rate during the Holocene remains to be determined in much of the Eastern Mediterranean.

Coastal Erosion in Libya

As in Gaza, erosion along the Cyrenaica coast (eastern Libya) has damaged archaeological sites, notably the Hellenistic and Roman harbor cities of Apollonia and Tocra, but also many smaller and less well-studied ancient sites (Bennett et al. 2004; Nikolaus, Abdrbba, and Emrage 2022; fig. 4). Several natural factors conspire to make this coastal stretch susceptible to erosion: extensive deposits of unconsolidated sediment close to the shore, a largely open coastline exposed to some of the highest waves in North Africa (Barbariol et al. 2021), and tectonic subsidence within the last two thousand years that has pulled archaeological structures originally built above high water closer to the water's edge (Flemming 2021).

Recent and twentieth-century shoreline movements have been quantified using satellite imagery, historic aerial photos, and declassified 1970s spy satellite (KH-9) images (Westley et al. 2023). Erosion rates of greater than two meters per year are present in the vicinity of Ptolemais, Tocra, and Apollonia, as well as rates of up to one-half meter per year in locations where archaeological material is either built on or buried within unconsolidated backshore sediments (fig. 5). This analysis also suggests that erosion rates have accelerated within the last ten-to-twenty years. If such rates had prevailed since the abandonment of these settlements, it is unlikely that many of the visible structures would still be preserved, given that these are ancient harbors with structures built close to the water's edge. This is exemplified at Apollonia where an 1828 historic map shows a broadly similar shoreline, albeit with localized instances of retreat. If erosion rates of two-to-five-tenths meter per year (as observed recently) had operated since 1820, this would have resulted in forty-to-one-hundred meters of coastal retreat, which does not seem to have been the case (fig. 6). What, therefore, could have caused this acceleration?

Comparison of KH-9 images from 1974 with recent images highlights how actively eroding areas were once fronted by extensive beaches that no longer exist. Without the natural protection afforded by these beaches, waves can reach the vulnerable unconsolidated backshore during storms and, possibly, during moderate or fair-weather conditions. Though not all the drivers of this land loss are known, one probable culprit is unregulated sand mining. This is suggested by satellite images and confirmed by local reports for the coast west of Ptolemais where erosion rates are highest (Elghazali 2019; fig. 7). Another possible factor is disruption of wadi-supplied sediment by urbanization and infrastructure that interferes with natural water flow, for example, roads, bridges, buildings, and culverts. Such features are evident from comparison of KH-9 and recent



satellite images (fig. 8). These observations tally with wider studies from North Africa that link widespread twentieth-century beach loss and coastal retreat to human factors operating independently of climate change, such as upstream damming and urbanization (Hzami et al. 2021).

Tropical Cyclones and Flooding in Oman

In recent years, countries in the Arabian Peninsula, particularly Yemen and Oman, have experienced the catastrophic impact of tropical cyclones on an almost yearly basis. Many have associated the increasing frequency and intensity of cyclones with climate change (Knutson et al. 2010; Al-Awadhi, Charabi, and Choudri 2019; Wehner, Zarzycki, and Patricola 2019).

The resulting coastal flooding is sufficiently serious that a warning system has been installed in Oman to mitigate future impacts. References to cyclone frequency that inform future projections rely on existing documentation by the National Oceanic and Atmospheric Administration (Knapp et al. 2010), which has recorded thirty-six “hurricanes” affecting the Omani coastline in the past 129 years (fig. 9). Though only hurricanes with sustained winds >150 mph are classified as tropical cyclones, hurricanes with lower intensity have had a catastrophic impact on Oman’s infrastructure and environment (Mansour et al. 2021).

Problematically, information on cyclone activity before the mid-twentieth century is scarce. Historical descriptions of strong winds that could classify as hurricanes or cyclones date to 1286 CE, 1325 CE,




Figure 9. Map of maritime archaeological sites in Oman documented in the EAMENA database in conjunction with major cyclone tracks over the past 130 years. Adapted from Andreou et al. 2022b: 142, fig. 9.

and 1853 CE (Newton and Zarins 2019), while historic flooding events (1600–1700 CE) have been identified in the sedimentary record near known sites, for example, at Al-Baleed in Dhofar (Reinhardt 2000)—the governorate with the highest exposure to tropical cyclones (fig. 10). Extreme sea level events have also been noted in the geoarchaeological record of Ras al-Hadd in northeastern Oman, dating to 4450 cal BP (Hoffman et al. 2015). Though the deposits at Ras al-Hadd are tentatively ascribed to a tsunami, there is a general understanding that more evidence for extreme wave events and flooding can and will be identified in future excavations.

Based on the available evidence, it appears that the precise relation between the increasing frequency and intensity of cyclones and climate change requires further historical, archaeological, and geological input to minimize any reasonable hypothesis that increasing frequency is the result of more effective documentation (e.g., Blount, Fritz, and Al-Harthy 2009). Similarly, further investigation is required to quantify the contribution of extensive construction of modern impervious surfaces (Al-Awadhi et al. 2016, 2018), particularly in coastal urban centers in Oman, which locally enhance the impact of flooding events.

Considerable attention given to the impact of tropical cyclones on Oman’s urban infrastructure (Hereher 2020; Mansour et al. 2021) has opened the way for assessments of the condition of archaeological sites located within the cyclone tracks through satellite imagery analysis (Andreou et al. 2022b, 2022c). Importantly, the identification of site clusters that are highly vulnerable to cyclone activity, particularly in the northeast (Muscat, Ash Sharqiyah) and southwest (Dhofar), could enable exploration of the premodern frequency of such events (e.g., Hoffmann et al. 2015; Ermertz et al. 2019). New geoarchaeological information will not only elucidate the frequency and intensity of historic cyclones but also has the potential to inform risk management strategies. Though heritage perspectives are often incorporated in risk management strategies, these perspectives tend to serve as cautionary tales within the scope of protection and preservation. With archaeology uniquely placed to add much-needed historical depth to increase the accuracy of climate change projections (see also Wright 2016), there is scope for geoarchaeological studies not only to confirm, but also to inform existing vulnerability models, thereby contributing more actively to heritage policies.

Contextualizing Maritime Archaeology in Climate Change Research

Climate change involves a dense network of interdependencies, including cultural heritage and modern human societies. We examine these interdependencies by reference to the coastal zone and the commonly cited destructive impacts of flooding and erosion, both of which will be aggravated by sea-level rise and changing storm patterns (ICOMOS 2019). The discussed case studies show that contemporary erosion and flooding impacts on coastal heritage are not solely attributable to climate change impacts but also include a strong contribution from human action. More specifically, activities that restrict coastal sediment supply,




Figure 10. Coastal erosion affecting the beach in front of Medieval Al-Baleed, Dhofar Governorate, Oman (top). Yellow box indicates a wooden shipwreck on the beach, exposed by storm action (bottom). Photographs by Tom Vosmer, 2013.

including sand mining, construction, and damming, can aggravate erosion on local and regional scales. Similarly, the extent and impact of flooding is contingent on human action: reduction through artificial flood defenses but exacerbation via impervious surfaces created by urbanization. In short, many coastal changes we see today that damage or destroy coastal archaeological sites, in the MENA region at least, result from human actions related to urban and economic development. A direct link between these changes and climate-change-driven sea-level rise is, at present, less clear. However, as we show below, this link will probably strengthen over the twenty-first century. Nonetheless, we emphasize that this observation is specific to our study region. Other parts of the world show clearer and stronger links with climate change, for instance the vastly accelerated erosion of polar and subpolar coastlines due to permafrost melting.

Consideration of the above suggests three main implications. First, from an archaeological preservation standpoint, the headline action of reducing greenhouse gas emissions is not enough. It must accompany wider environmental regulation that prevents or limits activities that damage both archaeological sites and natural environments. Unfortunately, many beaches have already been pushed past their point of recovery (Cooper and Jackson 2019). Moreover, at least some of the major infrastructure responsible for coastal change will probably remain in place for the foreseeable future. For example, many large dams built in the twentieth century are coming to the theoretical lower limit of their planned lifespan (usually ~50–120 years). However, the choice of deconstruction versus repair or renewal is complex and dependent on cost, politics, and whether substitutes are available for benefits, such as hydroelectric power. Also, dam building continues across the MENA region, with a new crop coming online over the coming decades (Marchetti et al. 2019). Nor will sea-level rise and its linked destructive impacts completely stop within the twenty-first century even if carbon emissions are reduced (IPCC 2022). Even so, we note that the main acceleration of SLR will come post-2050 if carbon emissions remain high: There is a difference of approximately forty centimeters between low and high emissions scenarios by 2100 compared to approximately five centimeters by 2050 (IPCC 2022; Vousdoukas et al. 2018). Arguably therefore, any action that can reduce exacerbating processes should be taken before this acceleration. These could include, for example, local regulation against destructive sand mining or more active consideration of the wider impacts of coastal hardening (e.g., Bitan et al. 2020). To be fully cognizant of archaeological management requirements, this necessitates maritime heritage to be integrated within wider coastal and marine management policies, something that is not standard practice, and that has been proposed in the context of marine protected areas (Breen et al. 2021).

Second, although large-scale models can project with reasonable certainty general patterns of sea-level rise, the unknown impact of unpredictable human activities makes it challenging to project twenty-first-century land-loss at smaller scales, such as the extent of an archaeological site or cluster of sites. Estimates are possible, particularly if they can be constrained by high spatial and temporal resolution measurement of local

trends in shoreline movement (Westley et al. 2023). But archaeologists, managers, and other stakeholders should always be prepared to assess critically model inputs, construction, and outputs; to consider a range of scenarios based on the drivers of change at a given site; and finally, to be aware that the certainty of these projections decreases the further forward in time one looks. This requires the expansion and diversification of existing, already interdisciplinary archaeological training to include quantitative skills necessary for assessing climate-change projections and their relevance to heritage management.

Third, big archaeological databases have allowed identification of large-scale patterns of landscape alterations at times linked with the broader impact of climate change (Westley et al. 2021). These databases offer significant opportunities in identifying and conducting in-depth geological, geomorphological, and paleoenvironmental research that will shed light on the frequency and intensity of phenomena attributed to climate change. Of course, identified patterns cannot be viewed outside of their unique economic and sociopolitical circumstances, the active consideration of which can be achieved through the incorporation of local communities (Stahl 2020) that have a daily engagement with their endangered heritage and who will ultimately need to respond to the impact of climate change on their heritage and environment.

The challenge of untangling multiple drivers of change in order to reach confident projections for the future is as important to maritime cultural heritage management as in other sectors. To prioritize action and to determine the most effective mitigation strategies in each context, archaeological practitioners and heritage managers require a clear idea of the nature, extent, and intensity of future impacts. Current global-scale models go partway to address this. They are useful from a broad-scale standpoint, mainly in terms of raising awareness of potential threat and addressing regional patterns. However, at a practical management level more detail is needed on individual sites. This is especially relevant in the coastal zones where past, present, and future impacts involve complex interactions of climate-change-driven processes and human action. There are clear challenges to achieving this in the Middle East and North Africa, notably in terms of the availability of high-resolution data at a frequency that can be used to track and quantify shoreline changes. This is slowly starting to change, particularly with increasing availability of satellite and UAV imagery, but in many cases the baseline datasets are relatively recent in time, thus making it harder to determine if identified trends are anomalous or reflective of long-term patterns.

We also note that those impacts highlighted here are but a subset of the full range of coastal complexity, which includes, for example, subsidence (enhanced by hydrocarbon and groundwater extraction), salinization (enhanced by fresh- and groundwater abstraction), seawater oxygen content (affected by pollution), and seabed erosion (enhanced by seagrass loss from pollution and bottom trawling; see also Gregory et al. 2022; ICOMOS 2019). Future changes will impact not only maritime archaeology but the natural environment and human societies. As such, there is a strong case that maritime archaeology should be better integrated into mitigation and management efforts that tend to focus

on the natural environment and modern infrastructure. Additionally, taking a long-term view and using (geo)archaeological evidence to extend the evidential baseline could contribute to the wider body of research that aims to disentangle the compounding processes driving change and, ultimately, assist in efforts to improve projections of future climate change impacts.

Acknowledgments

This work is part of the Maritime Endangered Archaeology Project (www.marea.soton.ac.uk) funded by Arcadia, a charitable fund of Lisbet Rausing and Peter Baldwin. Archaeological fieldwork in Libya was funded by the British Society for Libyan and North African Studies and the University of Southampton in collaboration with DoA Cyrenaica, Omar al-Mukhtar University, and Benghazi University. Fieldwork at Tell Ruqeish was funded by the Honor Frost Foundation and the University of Southampton, in collaboration with the Islamic University of Gaza and the Palestinian Ministry of Tourism. High resolution satellite images for eastern Libya and Gaza were provided by the European Space Agency.

References

- Abd-el Monsef, Hesham, Scot E. Smith, and Kamal Darwish. 2015. Impacts of the Aswan High Dam after 50 Years. *Water Resource Management* 29:1873–85. <https://doi.org/10.1007/s11269-015-0916-z>.
- Abualtayef, Mazen, Mohammed Abd Rabou, Samir Affi, Abdel Fattah Abd Rabou, Ahmed Khaled Seif, and Ali Masria. 2021. Change Detection of Gaza Coastal Zone Using GIS and Remote Sensing Techniques. *Journal of Coastal Conservation* 25:1–20. <https://doi.org/10.1007/s11852-021-00825-4>.
- Abualtayef, Mazen, Said Ghabayen, Ahmed Abu Foul, Ahmed Seif, Masamitsu Kuroiwa, Yuhei Matsubara, and Omar Matar. 2012. The Impact of Gaza Fishing Harbour on the Mediterranean Coast of Gaza. *Journal of Coastal Development* 16:1–10.
- Al-Awadhi, Talal, Yassine Charabi, and Bheemanagoud S. Choudri. 2019. Pathways for Building Urban Resilience to Climate Change in Oman. *Development in Practice* 29:594–605. <https://doi.org/10.1080/09614524.2019.1593320>.
- Al-Awadhi, Talal, Yassine Charabi, Bheemanagoud S. Choudri, and Y. Bani Oraba. 2018. Flooding Risk Analysis: A Case Study of Muscat Governorate, Sultanate of Oman. *Human and Ecological Risk Assessment: An International Journal* 24:667–78. <https://doi.org/10.1080/10807039.2017.1396441>.
- Al-Awadhi, Talal, Elnazir Ramadan, Bheemanagoud S. Choudri, and Yassine Charabi. 2016. Growth of Coastal Population: Likely Exposure to Sea-level Rise and Associated Storm Surge Flooding in the Sultanate of Oman. *Journal of Environmental Management and Tourism* 7:341–46.
- Anderson, David, Thaddeus Bissett, Stephen J. Yerka, Joshua J. Wells, Eric C. Kansa, Sarah W. Kansa, Kelsey Noack Myers, R. Carl DeMuth, and Devin A. White. 2017. Sea-Level Rise and Archaeological Site Destruction: An Example from the Southeastern United States Using DINAA (Digital Index of North American Archaeology). *PLoS ONE* 12:e0188142. <https://doi.org/10.1371/journal.pone.0188142>.
- Andreou, Georgia M., Michael Fradley, Lucy Blue, and Colin Breen. 2022a. Establishing a Baseline for the Study of Maritime Cultural Heritage in the Gaza Strip. *Palestine Exploration Quarterly*. <https://doi.org/10.1080/00310328.2022.2037923>.
- Andreou, Georgia M., Julia Nikolaus, Kieran Westley, Crystal El. Safadi, Lucy Blue, Ash Smith, and Colin Breen. 2022b. Big Data in Maritime Archaeology: Challenges and Prospects from the Middle East and North Africa. *Journal of Field Archaeology* 47:131–48. <https://doi.org/10.1080/00934690.2022.2028082>.
- Andreou, Georgia M., Kieran Westley, Harmen O. Huigens, and Lucy Blue. 2022c. Exploring the Impact of Tropical Cyclones on Oman's Maritime Cultural Heritage through the Lens of Al-Baleed, Salalah (Dhofar Governorate). *Journal of Maritime Archaeology* 17:465–86. <https://doi.org/10.1007/s11457-022-09333-4>.
- Anthony, Edward, Nick Marriner, and Christophe Morhange. 2014. Human Influence and the Changing Geomorphology of Mediterranean Deltas and Coasts Over the Last 6000 Years: From Progradation to Destruction Phase? *Earth Science Reviews* 139:336–61. <https://doi.org/10.1016/j.earscirev.2014.10.003>.
- Barbariol, Francesco, Silvio Davison, Francesco Marcello Falcieri, Rossella Ferretti, Antonio Ricchi, Mauro Scavo, and Alvis Benetazzo. 2021. Wind Waves in the Mediterranean Sea: An ERA5 Reanalysis Wind-Based Climatology. *Frontiers in Marine Science* 8:1615. <https://doi.org/10.3389/FMARS.2021.760614>.
- Barkai, Ofra, Oded Katz, Amit Muskhin, and Beverly Goodman-Tchernov. 2018. Long-Term Retreat Rates of Israel's Mediterranean Sea Cliffs Inferred from Reconstruction of Eroded Archaeological Sites. *Geoarchaeology* 33:314–27. <https://doi.org/10.1002/geo.21633>.
- Beechey, Frederick, and Henry Beechey. 1828. *Proceedings of the Expedition to Explore the Northern Coast of Africa*. London: Murray.
- Bennett, Paul, Andrew Wilson, Ahmed Buzaian, and Alette Kattenberg. 2004. The Effects of Recent Storms on the Exposed Coastline of Tobra. *Libyan Studies* 35:113–22. <https://doi.org/10.1017/S0263718900003745>.
- Bitan, Menashe, Ehud Galili, Ehud Spanier, and Dov Zviely. 2020. Beach Nourishment Alternatives for Mitigating Erosion of the Ancient Coastal Sites on the Mediterranean Coast of Israel. *Journal of Marine Science and Engineering* 8:509. <https://doi.org/10.3390/jmse8070509>.
- Blount, Christopher, Hermann Fritz, and Ahmed Hamoud Mohammed Al-Harthy. 2009. Coastal Vulnerability Assessment Based on Historic Tropical Cyclones in the Arabian Sea. Pp. 207–14 in *Indian Ocean Tropical Cyclones and Climate Change*, ed. Yassine Charabi and Salim Al-Hatrushi. Dordrecht: Springer. http://dx.doi.org/10.1007/978-90-481-3109-9_25.
- Breen, Colin, Crystal El Safadi, Harmen O. Huigens, Sophie Tews,

- Kieran Westley, Georgia Andreou, Rodrigo Ortiz Vazquez, Julia Nikolaus, and Lucy Blue. 2021. Integrating Cultural and Natural Heritage Approaches to Marine Protected Areas in the MENA Region. *Marine Policy* 132:104676. <https://doi.org/10.1016/j.marpol.2021.104676>.
- Brooks, Nick, Joanna Clarke, Grace Wambui Ngaruiya, and Elizabeth Edna Wangui. 2020. African Heritage in a Changing Climate. *Azania* 55:287–328. <https://doi.org/10.1080/0067270X.2020.1792177>.
- Cooper, J. Andrew G., and Derek W. T. Jackson. 2019. Coasts in Peril? A Shoreline Health Perspective. *Frontiers in Earth Science* 7. <https://doi.org/10.3389/feart.2019.00260>.
- Cooper, J. Andrew G., Gerd Masselink, Gianluca Coco, Andy D. Short, Bruno Castelle, Kerry Lee Rogers, Edward Anthony, Andrew N. Green, Joseph T. Kelley, Orrin H. Pilkey, and Derek W. T. Jackson. 2020. Sandy Beaches Can Survive Sea-Level Rise. *Nature Climate Change* 10:993–95. <https://doi.org/10.1038/s41558-020-00934-2>.
- Dawson, Tom, Joanna Hambly, William Lees, and Sara Miller. 2021. Proposed Policy Guidelines for Managing Heritage at Risk Based on Public Engagement and Communicating Climate Change. *The Historic Environment: Policy & Practice* 12:375–394. <https://doi.org/10.1080/17567505.2021.1963573>.
- Derooin, Jean-Paul, Rania Bou Kheir, and Chadi Abdullah. 2017. Geoarchaeological Remote Sensing Survey for Cultural Heritage Management: Case Study from Byblos (Jbail, Lebanon). *Journal of Cultural Heritage* 23:37–43. <https://doi.org/10.1016/j.culher.2016.04.014>.
- Elgazali, Isra. 2019. Environmental Impact of Sand Mining on the Vegetation and Surrounding Environment in the Coast of Botrabah, the Eastern Coast of Libya (Cyrenaeca) Coastal Erosion. BS thesis, University of Benghazi.
- Ermert, Alina Marie, Miklos Kázmér, Silja Kerstin Adolphs, Michaela Falkenroth, and Gösta Hoffmann. 2019. Geoarchaeological Evidence for the Decline of the Medieval City of Qalhat, Oman. *Open Quaternary* 5:8. <https://doi.org/10.5334/oq.56>.
- Flemming, Nicholas. 2021. *Apollonia on My Mind: The Memoir of a Paraplegic Ocean Scientist*. Leiden: Sidestone Press.
- Galili, Ehud, and Baruch Rosen. 2010. Preserving the Maritime Cultural Heritage of the Mediterranean, a Cradle Of Cultures, Religions and Civilizations—The Holy Land Perspective. *Journal of Coastal Conservation* 14:303–15. <https://doi.org/10.1007/s11852-010-0107-2>.
- Gregory, David, Tom Dawson, Dolores Elkin, Hans van Tilburg, Chris Underwood, Vicki Richards, Andrew Viduka, Kieran Westley, Jeneva Wright, and Jørgen Hollesen. 2022. Of Time and Tide: The Complex Impacts of Climate Change on Coastal and Underwater Cultural Heritage. *Antiquity* 96.390:1396–411. <https://doi.org/10.15184/aqy.2022.115>.
- Hambrecht, George, and Marcy Rockman. 2017. International Approaches to Climate Change and Cultural Heritage. *American Antiquity* 82:627–41. <https://doi.org/10.1017/aaq.2017.30>.
- Hereher, Mohamed E. 2020. Assessment of Infrastructure Vulnerability to Tsunamis upon the Coastal Zone of Oman Using GIS. *Geosciences* 10:175. <https://doi.org/10.3390/geosciences10050175>.
- Hinkel, Jochen, Luc Feyen, Mark A. Hemer, Gonéri Le Cozannet, Daniel Lincke, Marta Marcos, Lorenzo Mentaschi Jan L. Merckens, Hans de Moel, Sanne Muis, Robert J. Nicholls, Athanasios T. Vafeidis, Roderik S. W. van de Wal, Michalis I. Vousdoukas, Thomas Wahl, Philip J. Ward, and Claudia Wolff. 2021. Uncertainty and Bias in Global to Regional Scale Assessments of Current and Future Coastal Flood Risk. *Earth's Future* 9:e2020EF001882. <https://doi.org/10.1029/2020EF001882>.
- Hoffman, Gösta., Christoph Grützner, Klaus Reicherter, and Frank Preusser. 2015. Geo-Archaeological Evidence for a Holocene Extreme Flooding Event within the Arabian Sea (Ras al Hadd, Oman). *Quaternary Science Reviews* 113:1123–33. <https://doi.org/10.1016/j.quascirev.2014.09.033>.
- Hzami, Abderraouf, Essam Heggy, Oula Amrouni, Gil Mahé, Mohamed Maanan, and Saâdi Abdeljaouad. 2021. Alarming Coastal Vulnerability of the Deltaic and Sandy Beaches of North Africa. *Scientific Reports* 11:2320. <https://doi.org/10.1038/s41598-020-77926-x>.
- ICOMOS. 2019. *The Future of Our Pasts: Engaging Cultural Heritage in Climate Action*. Technical Report. Paris: ICOMOS. <https://civvih.icomos.org/wp-content/uploads/Future-of-Our-Pasts-Report-min.pdf>.
- IPCC. 2022. *IPCC Special Report on the Ocean and Cryosphere in a Changing Climate*, ed. Hans-Otto Portner et al. Cambridge: Cambridge University Press. <https://doi.org/10.1017/9781009157964>.
- Klinger, Philip, and Kim Knauer. 2020. Shoreline Mapping in the Gaza Strip. https://eo4society.esa.int/wp-content/uploads/2020/10/EOC0011_WOR_v01.pdf.
- Knapp, Kenneth R., Michael C. Kruk, David H. Levinson, Howard J. Diamond, and Charles J. Neuman. 2010. The International Best Track Archive for Climate Stewardship (IBTraCS): Unifying Tropical Cyclone Data. *Bulletin of the American Meteorological Society* 91:363–76. <https://doi.org/10.1175/2009BAMS2755.1>.
- Knutson, Thomas R., John L. McBride, Johnny Chan, Kerry Emanuel, Greg Holland, Chris Landsea, Isaac Held, James P. Kossin, A. K. Srivastana, and Masato Sugi. 2010. Tropical Cyclones and Climate Change. *Nature Geoscience* 3:157–63. <https://doi.org/10.1038/ngeo779>.
- Li, Yuqi, Xin Jia, Zhen Liu, Luo Zhao, Pengfei Sheng, and Michael J. Storzum. 2022. The Potential Impact of Rising Sea Levels on China's Coastal Cultural Heritage: A GIS Risk Assessment. *Antiquity* 96.386:406–21. <https://doi.org/10.15184/aqy.2022.1>.
- Mansour, Shawky, Stephen Darby, Julian Leyland, and Peter M. Atkinson. 2021. Geospatial Modelling of Tropical Cyclone Risk along the Northeast Coast of Oman: Marine Hazard Mitigation and Management Policies. *Marine Policy* 129:104544. <https://doi.org/10.1016/j.marpol.2021.104544>.
- Marchetti, Nicolò, Antonio Curci, Maria Carmela Gatto, Serena Nicolini, Simone Mühl, and Federico Zaina. 2019.

- A Multi-Scalar Approach for Assessing the Impact of Dams on the Cultural Heritage in the Middle East and North Africa. *Journal of Cultural Heritage* 37:17–28. <https://doi.org/10.1016/j.culher.2018.10.007>.
- Marriner, Nick, Jean-Philippe Goiran, Bernard Geyer, Valérie Matoïan, Michel al-Maqdissi, Marion Leconte, and Pierre Carbonel. 2012. Ancient Harbors and Holocene Morphogenesis of the Ras Ibn Hani Peninsula (Syria). *Quaternary Research* 78:35–49. <https://doi.org/10.1016/j.yqres.2012.03.005>.
- Newton, Lynne, and Juris Zarins. 2019. *Dhofar through the Ages: An Ecological, Archaeological and Historical Landscape*. Archaeological Heritage of Oman. Oxford: Archaeopress. <https://doi.org/10.2307/j.ctvndv7n0>.
- Nikolaus, Julia, Mohamed O. M. Abdrrbba, and Ahmad Emrage. 2022. The Cyrenaica Coastal Survey Project: Documenting Endangered Maritime Heritage in Libya. *Journal of Maritime Archaeology* 17:421–44. <https://doi.org/10.1007/s11457-022-09336-1>.
- Reimann, Lena, Athanasios T. Vafeidis, Sally Brown, Jochen Hinkel, and Richard S. J. Tol. 2018. Mediterranean UNESCO World Heritage at Risk from Coastal Flooding and Erosion Due to Sea-Level Rise. *Nature Communications* 9:4161. <https://doi.org/10.1038/s41467-018-06645-9>.
- Reinhardt, Eduard. 2000. Evolution of the Ancient Harbor at Al Balid, Oman: Sedimentological and Micropalaeontological Evidence. PhD diss., McMaster University.
- Sesana, Eleana, Alexandre S. Gagnon, Chiara Ciantelli, JoAnn Casar, and John J. Hughes. 2021. Climate Change Impacts on Cultural Heritage: A Literature Review. *WIREs Climate Change* 12:e710. <https://doi.org/10.1002/wcc.710>.
- Stahl, Ann Brower. 2020. Assembling “Effective Archaeologies” toward Equitable Futures. *American Anthropologist* 122:37–50. <https://doi.org/10.1111/aman.13365>.
- Vousdoukas, Michalis I., Joanne Clarke, Roshanka Ranasinghe, Lena Reimann, Nadia Khalaf, Trang Minh Duong, Birgitt Ouweneel, Salma Sabour, Carley E. Iles, Christopher H. Trisos, Luc Feyen, Lorenzo Mentaschi, and Nicholas P. Simpson. 2022. African Heritage Sites Threatened as Sea-Level Rise Accelerates. *Nature Climate Change* 12:256–62. <https://doi.org/10.1038/s41558-022-01280-1>.
- Vousdoukas, Michalis I., Lorenzo Mentaschi, Evangelos Voukoulalas, Martin Verlaan, Svetlana Jevrejeva, Luke P. Jackson, and Luc Feyen. 2018. Global Probabilistic Projections of Extreme Sea Levels Show Intensification of Coastal Flood Hazard. *Nature Communications* 9:2360. <https://doi.org/10.1038/s41467-018-04692-w>.
- Vousdoukas, Michalis I., Roshanka Ranasinghe, Lorenzo Mentaschi, Theocharis A. Plomaritis, Panagiotis Athanasiou, Arjen Luijendijk, and Luc Feyen. 2020. Sandy Coastlines under Threat of Erosion. *Nature Climate Change* 10:260–63. <https://doi.org/10.1038/s41558-020-0697-0>.
- Wehner, Michael, Colin Zarzycki, and Christina Patricola. 2019. Estimating the Human Influence on Tropical Cyclone Intensity as the Climate Changes. Pp.235–60 in *Hurricane Risk* ed. Jennifer M. Collins, and Kevin Walsh. Cham: Springer. https://doi.org/10.1007/978-3-030-02402-4_12.
- Westley, Kieran, Georgia Andreou, Crystal El Safadi, Harmen O. Huigens, Julia Nikolaus, Rodrigo Ortiz-Vazquez, Nick Ray, Ash Smith, Sophie Tews, Lucy Blue, and Colin Breen. 2021. Climate Change and Coastal Archaeology in the Middle East and North Africa: Assessing Past Impacts and Future Threats. *Journal of Island and Coastal Archaeology* 18:251–83. <https://doi.org/10.1080/15564894.2021.1955778>.
- Westley, Kieran, Julia Nikolaus, Ahmed Emrage, Nic Flemming, and Andrew Cooper. 2023. The Impact of Coastal Erosion on the Archaeology of the Cyrenaican Coast of Eastern Libya. *PLOS ONE* 18:e0283703. <https://doi.org/10.1371/journal.pone.0283703>.
- Williams, Allan T., Nelson Rangel-Buitrago, Enzo Pranzini, and Giorgio Anfuso. 2018. The Management of Coastal Erosion. *Ocean & Coastal Management* 156:4–20. <https://doi.org/10.1016/j.ocecoaman.2017.03.022>.
- Wright, Jeneva. 2016. Maritime Archaeology and Climate Change: An Invitation. *Journal of Maritime Archaeology* 11:255–70. <https://doi.org/10.1007/s11457-016-9164-5>.
- Yasur-Landau, Assaf, Gilad Shtienberg, Gil Gambash, Giorgio Spada, Daniele Melini, Ehud Arkin-Shalev, Anthony Tamberino, Jack Reese, Thomas E. Levy, and Dorit Sivan. 2021. New Relative Sea-Level (RSL) Indications from the Eastern Mediterranean: Middle Bronze Age to the Roman Period (~3800–1800 y BP), Archaeological Constructions at Dor, the Carmel Coast, Israel. *PLOS ONE* 16:e0251870. <https://doi.org/10.1371/journal.pone.0251870>.



Kieran Westley is a research fellow at the School of Geography & Environmental Science, Ulster University (UK). He is currently employed on the Maritime Endangered Archaeology (MarEA) project where he is researching climate change impacts and threats to maritime archaeology in the Middle East and North Africa. He has wider research interests in submerged prehistoric landscapes and the application of marine and satellite remote sensing to archaeological research.

Georgia Andreou is a research associate at the Maritime Endangered Archaeology Project at the University of Southampton and an associate lecturer in Near Eastern Archaeology at UCL. Her research focuses on the second and first millennia BCE in the eastern Mediterranean with emphasis on coastal archaeological sites. She is currently directing the Gaza Maritime Archaeology Project (GAZAMAP). She is also one of the codirectors of the Kalavassos and Maroni Built Environments Project (KAMBE) in Cyprus.

