


 The Holocene

**Environment-man relationships in historical times: the
balance between urban development and natural forces at
Leptis Magna (Libya)**

Journal:	<i>The Holocene</i>
Manuscript ID:	Draft
Manuscript Type:	Paper
Date Submitted by the Author:	
Complete List of Authors:	Pucci, Stefano; Istituto Nazionale di Geofisica e Vulcanologia Pantosti, Daniela; Istituto Nazionale di Geofisica e Vulcanologia De Martini, Paolo Marco; Istituto Nazionale di Geofisica e Vulcanologia Smedile, Alessandra; Istituto Nazionale di Geofisica e Vulcanologia Munzi, Massimiliano; Comune di Roma, Sovraintendenza ai Beni Culturali Cirelli, Enrico; Università di Bologna, Dipartimento di Archeologia Pentiricci, Massimo; Comune di Roma, Sovraintendenza ai Beni Culturali Musso, Luisa; Università degli studi Roma3
Keywords:	natural hazards, geomorphology, streams dynamics, quaternary

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Abstract:	<p>We analyze the relationships between human modification of the environment and natural events in the Roman city of Leptis Magna (UNESCO world heritage), western Libya. For the first time the history of Leptis Magna is tested against a high resolution geomorphological and stratigraphical reconstruction and absolute radiocarbon dating.</p> <p>Historical and archaeological sources indicate the occurrence of different natural events as the cause of the town decline: earthquakes, flooding, and tsunamis. Our data indicate that Leptis Magna was tightly linked to the nearby wadi. In fact, aware of the hazards related to violent flooding, the Romans were able to cope with the threat posed by the wadi by engineering defensive hydraulic works around the town. Once economic decay began and the society could no longer guarantee the ongoing maintenance of these structures, the decline of the settlement began and the occurrence of violent floods reclaimed the populated areas. The occurrence of large earthquakes, and the tsunami event that damaged the harbor have been discarded as playing a primary role in the decline of Leptis Magna.</p>
-----------	---



1
2
3 **Environment-man relationships in historical times: the balance between urban**
4 **development and natural forces at Leptis Magna (Libya).**
5
6
7
8
9

10 **Pucci, S.^{1*}, D. Pantosti¹, P.M. De Martini¹, A. Smedile¹, M. Munzi², E. Cirelli³, M.**
11 **Pentiricci², L. Musso⁴.**
12
13
14
15
16

17 ¹ Istituto Nazionale di Geofisica e Vulcanologia, Via di Vigna Murata, 605, 00143 Roma, Italy

18 ² Sovraintendenza ai Beni Culturali del Comune di Roma, Via del Teatro Marcello 5, 00186 Roma, Italy

19 ³ Dipartimento di Archeologia, Università di Bologna, Via San Vitale 28, 48100 Bologna, Italy

20 ⁴ Università degli studi Roma3, Piazza della Repubblica, 10, 00185 Roma, Italy

21 * Corresponding Author: pucci@ingv.it
22
23
24
25
26
27
28
29
30
31

32 **Abstract**

33
34 We analyze the relationships between human modification of the environment and natural
35 events in the Roman city of Leptis Magna (UNESCO world heritage), western Libya. For the first
36 time the history of Leptis Magna is tested against a high resolution geo-morphological and
37 stratigraphical reconstruction and absolute radiocarbon dating.
38
39
40
41
42

43 Historical and archaeological sources indicate the occurrence of different natural events as
44 the cause of the town decline: earthquakes, flooding, and tsunamis. Our data indicate that Leptis
45 Magna was tightly linked to the nearby wadi. In fact, aware of the hazards related to violent
46 flooding, the Romans were able to cope with the threat posed by the wadi by engineering defensive
47 hydraulic works around the town. Once economic decay began and the society could no longer
48 guarantee the ongoing maintenance of these structures, the decline of the settlement began and the
49 occurrence of violent floods reclaimed the populated areas. The occurrence of large earthquakes,
50 and the tsunami event that damaged the harbor have been discarded as playing a primary role in the
51 decline of Leptis Magna.
52
53
54
55
56
57
58
59
60

Introduction

Since ancient times, socio-economic strategies of some civilizations have led to undertake severe environmental modifications, that in some cases may become so drastic as to require natural risk management. On the other hand, the rise and fall of great civilizations, wars and human achievements, thus human history in general has been tightly related to extreme natural disasters. Societies responded to natural disasters according to their socio-economic state: in some cases a society was resilient to perturbations and in others a society was so vulnerable to perturbations (generally for economic reasons) that it was unable to cope and collapsed, failed or migrated elsewhere. Historical disasters left permanent marks on the cultural development of entire regions. These are recognized not only in the study of civilizations, urbanization, migrations etc., but also in the geology and geomorphology of an area.

The sensitivity or vulnerability of landscapes and ecosystems to human activities were critical issues even during the Roman Empire. In this paper we analyze human modification on the natural environment and, *vice versa*, of the natural events on human history. We focus on the city of Leptis Magna, a magnificent Roman town in Tripolitania (western Libya) that reached its maximum expansion during the Empire of Septimius Severus (193-211 A.D.) and that started to decline during the late IV century.

Leptis Magna is located on the North-African coastline at the outlet of a major dry-land stream (also called wadi or oued) that served as a natural harbor. This natural harbor was progressively enlarged starting with the work of Nero and reached its maximum development during the time of Septimius Severus. Historical and archaeological sources suggest that the harbor basin became inefficient and was abandoned due to its complete infill, possibly related to: i) violent flooding following the collapse of a dam built to regulate the course of the wadi because of the large 365 A.D. earthquake (Salza Prina Ricotti, 1995; Di Vita, 1990); ii) lack of maintenance due to the decline of the settlement induced by severe damage after the 365 A.D. earthquake (Di Vita, 1990

1
2
3 and 1995), or other local seismic sources (Guidoboni *et al.*, 1994; Stiros, 2001); iii) inundation of a
4
5 tsunami wave caused by the 365 A.D. earthquake that left a huge amount of debris and modified the
6
7 local coastal morphology (Guidoboni *et al.*, 1994; Ambraseys *et al.*, 1994; Lorito *et al.*, 2007, Shaw
8
9 *et al.*, 2008); iv) bad orientation and geometry of the harbor structures with respect to the local
10
11 marine currents that were bringing debris inside (Salza Prina Ricotti, 1972-1973).
12
13
14

15 In this work we present geological data, supported by geomorphological analysis and
16
17 radiocarbon dating, with the aim to: 1) verify the hypotheses concerning the harbor abandonment;
18
19 2) reconstruct the main events of the close human-environment interaction; 3) discover if there is a
20
21 cause-effect relationship between natural disasters and the settlement decline; 4) integrate the
22
23 archaeological knowledge at Leptis Magna by providing some independent chronological
24
25 constraints.
26
27
28
29
30
31
32
33

34 **Methodology**

35
36 The methodology used here is based on a geological and geomorphological survey of the
37
38 area in order to understand the depositional and erosional systems driven by the dynamics of the
39
40 wadi. Drainage is certainly the most sensitive system to external modifications such as those
41
42 induced by climate and human activity. As characteristic geomorphic markers of the wadi response
43
44 to such modifications, we reconstructed and mapped the flights of inset terrace surfaces along the
45
46 main channel of the Wadi Lebda. These terraces provide insights in to the degradational behavior of
47
48 the wadi following alluvial phases. For this purpose, a topographical survey of the study site was
49
50 performed by means of a differential GPS instrument (Leica Geosystems 1230), with profiles of 1.0
51
52 m horizontal resolution and 0.5 m maximum vertical standard deviation. Additionally, we
53
54 investigate the alluvial phases along the main channel and at the wadi outlet by applying traditional
55
56 stratigraphic analyses of outcrops and hand-made cores (up to 4.0 m deep) integrated with
57
58 micropaleontological and paleoenvironmental analysis. Depositional characteristics were examined
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

and correlated through the study area with the aim of recognizing changes to the wadi environment during historical times.

Because of the strict interaction of natural processes with human life, the stratigraphy included successive archaeological occupational layers. The stratigraphic position of archaeological artifacts (i.e. Roman villas and cisterns, Arabic fountains, etc.) and dating of ceramic artifacts were used as further constraints to complete this reconstruction.

An absolute chronological reference was provided by radiocarbon dating. We dated a total of 7 small fragments of charcoal sampled in selected layers at the Poznań Radiocarbon Laboratory (Poland). The samples were pre-treated with the acid-alkali-acid method and dated by decay counting through accelerator mass spectrometry (AMS) according to Goslar *et al.* (2004). The measured ages were dendrochronologically corrected for $^{12}\text{C}/^{14}\text{C}$ variations in the atmosphere through the radiocarbon calibration program Calib Rev 5.0.211 (Stuiver and Reimer, 1993) (Table I). The calibrated ages are calculated for the probability density function at the 95.4% (2σ) and at 68.3% (1σ) confidence limits. For the Roman Dam dating, seriating data were statistically treated through both combination and sequence models of Oxford Research Lab calibration program Oxcal 4.0. Because the dated samples are small fragments of charcoal, they could have experienced some reworking before deposition in hosting the sediment. Therefore, the possibility that the age of the sample could be older than the actual age of the sediment should be considered and represents an uncertainty.

The micropaleontological analysis consisted of quantitative and qualitative studies of the benthic foraminifera assemblages. Samples containing well preserved and abundant benthic foraminifera were counted (at least 100 specimens) and identified in the size fraction $< 125 \mu\text{m}$.

Further investigations on some selected samples were carried out through a cross sectional scanning electron microscopy (FESEM) analysis in order to discriminate the mineralogical content and to elucidate the possible relationships between different samples.

The Town

Since Phoenician times, the location of Leptis Magna was favorable to settlement because it offered both the availability of fresh water and a natural harbor that was strategically important for Mediterranean trade routes. After the Second Punic War, in the II c. B.C., the control of Leptis shifted from Carthage to Rome (Sallust, 40 B.C.). The city obtained the status of Municipium in the Flavian age (74-77 A.D.) and became a Colony during the Empire of Trajan (109 A.D.). This magnificent town continued to be enriched with monumental buildings between the I and the II c. A.D. Since it was his birthplace, the Emperor Septimius Severus led the town through its flourishing period, as testified by the proliferation of public monuments (the enlargement of the harbor, a great *nymphaeum*, a new *forum* with a magnificent basilica, etc.) and its general expansion, housing at its peak up to tens of thousands of citizens. The decline of the Roman town, commencing between the end of the IV and the beginning of the V c. A.D., was followed by a progressive contraction of the settlement during the Vandal (439-533 A.D.), Byzantine (533-644 A.D.) and Arabic times (following the conquest in 644 A.D.).

With the aim of understanding the depositional environment and the timing of depositional events, we studied five stratigraphic sections of the deposits that overlie the town. The section near the Septimius Severus Arch (S1, Fig. 1), along the *Cardo* (the main street), exposes a 4 m-thick, clear sequence of flooding events (Fig. 2a). Detailed analysis shows the lower 0.8 m of the alluvial deposit to be formed by a multilayer consisting of planar, parallel and laminated bedding of fine to coarse sand in a residual silty matrix. Local horizons of imbricated anthropic detritus (stones and pottery) cover the base of the ruins. Conversely, the upper part is characterized by massive deposits of a homogeneous, loose, fine to medium, mature quartzitic sand within a residual silty red matrix containing a few badly preserved (likely reworked) benthonic foraminifera. Towards the top, bioturbation occurs that conceals the original stratigraphy. Soil cover is absent or very thin

1
2
3 indicating young sediments and unfavorable conditions for pedogenesis, perhaps due to frequent
4 events of deposition/erosion. These deposits are equivalent to those observed in the wadi channels
5 and derive from the reworking of the Red Sand and Marine Quaternary formations outcropping in
6 the hinterland (Floridia, 1939; Lipparini, 1940; Capitanio, 1999). The characteristics of these
7 polycyclic deposits, the gentle (ca. 0.5°) seaward dip of bedding (Fig. 2b), along with the slightly
8 convex surface morphology of the alluvium, suggests that Leptis Magna is located on an alluvial
9 plain built up mainly from the nearby Wadi Lebda; thus, in an area prone to inundation. When the
10 town was excavated at the beginning of the 19th's, all its magnificence was revealed by removal of
11 alluvial deposits and sand derived from the active sand dunes that locally overlie the alluvium. Even
12 today it is possible to draw a flat surface across the ruins that represents the last alluvial surface
13 predating the archaeological excavations (Fig. 2c).
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28

29 This confirms that the main role in the town-environment interaction is played by the local
30 drainage network and its major dry-land stream, the Wadi Lebda. In semiarid Mediterranean
31 environments rock and soil erosion constitutes a major aspect of landscape degradation. In fact,
32 wadis discharge through quick, violent events (Schick, 1977; Reid and Laronne, 1995; Serrat *et al.*,
33 2001). Their suspended sediment yield is highly variable because precipitation and runoff are
34 themselves highly variable, with high sediment transport efficiency according to intermittent flow
35 regimes (Reid and Laronne, 1995). This highly erosive power has major repercussions both in the
36 eroded zone and in the receiving zone (inundations, silting up of channels, reservoirs and harbors).
37
38
39
40
41
42
43
44
45
46
47

48 Despite the possibility of the occurrence of large flooding events, both the Greeks and
49 Romans continued their occupation of the site because of its strategic location. Urban expansion
50 during Roman times occurred closer and closer to the outlet of the Wadi Lebda and was adjusted
51 following specific risk management plans. This flourishing society developed important hydraulic
52 and engineering works, not only to provide water to the town and its numerous baths or to enlarge
53 its trade capability, but especially to protect the town and its infrastructures from large and
54 damaging floods.
55
56
57
58
59
60

1
2
3 A sample collected at the base of section S1, thus dating the oldest flood deposit, was
4 radiocarbon dated at 320-440 A.D. (Table I). This age range indicates the beginning of exceptional
5 alluviation episodes affecting the urban area. Because these deposits were not removed, as expected
6 if there was systematic maintenance of the town, this suggests the beginning of the decline of
7 Leptis, when men gave up against natural hazards.
8
9
10
11
12
13
14
15
16
17
18
19

20 **The *agger* and the dam**

21
22 In order to protect Leptis Magna and its agricultural suburbs from potential aggressors, and
23 to keep away hazards of flooding, especially due to the presence of urban buildings along the wadi,
24 **the Romans erected a barrier of dirt, earth and soil (*agger*), up to 5.0 m high, with a ditch at its base.**
25 **According to Rinaldi Tufi (2000), this *agger* was built initially in 69 A.D.** between the town and the
26 hinterland. Stratigraphical analysis of the *agger* confirms its man-made nature: chaotic arrangement
27 of the alluvial deposits (i.e. loose, fine to medium sand and silt, with reworked benthonic
28 foraminifera, within residual silty red matrix) derived from excavation of the ditch, mixed with
29 artifacts and accumulation of skeletonized human bones. The abundance of well preserved charcoal
30 in the *agger* allowed its construction to be dated between 0 and 140 A.D. (S2, Fig. 1 and Table I).
31
32
33
34
35
36
37
38
39
40
41
42

43 The *agger* surrounds the town for several kilometers and connects with a stonework dam
44 where it crosses the Wadi Lebda. **The dam served not only as a water reservoir but also as a**
45 **strategic structure to regiment and deviate the flow of the Wadi Lebda along the *agger* into a**
46 **secondary wadi (er-Rsaf) to the west** (Fig. 1). The age of construction of the dam is not known, but
47 most authors suggest that it was built during the Empire of Septimius Severus to allow the
48 construction and functioning of the monumental harbor (Di Vita, 1990). The dam is 220 m long and
49 6 m high, the central part of which is made of thick walls with 4 ramparts. The dam body is still
50 standing and does not show any evidence of damage related to shaking induced by earthquakes.
51
52
53
54
55
56
57
58
59
60
Nowadays, the Wadi Lebda flows where the eastern shoulder of the dam was located, down-cutting

1
2
3 and under-excavating the artifact and further endangering its stability. **The whole basinward part of**
4 **the dam is buried by a depositional terrace for its entire height** (Fig. 3a). Geomorphological analysis
5
6 allowed us to recognize this terrace as formed by the reservoir siltation process, since no correlative
7
8 natural terraces along the wadi Lebda were identified (wadi terrace III, Fig. 1). In fact, the earliest
9
10 wadi bed predating the construction of the dam is still visible below the dam and coincides with the
11
12 central part of the construction (Fig. 3b). Nearby the dam shoulder, analysis of a stratigraphical
13
14 sequence (S3, Fig. 1) also showed pre-dam wadi deposits composed of laminated silt and fine sand
15
16 and interbedded poorly-developed paleosoils. The sequence shows well-stratified concave bedding
17
18 of a channel base, lying below the 4.5 m-thick sediments filling the reservoir, characterized by
19
20 homogeneous, compact silt and fine sand bodies, alternating with loose fine to medium sand, within
21
22 abundant residual silty red matrix (Fig. 3c). The channel base deposits were radiocarbon dated at 40
23
24 B.C.-90 A.D. (Table I), indicating that during this period the dam was not yet built. **Once the**
25
26 **Leptitans built the dam, the siltation was induced by the aggradational behavior of the Wadi Lebda,**
27
28 as a result of the modification of the natural longitudinal profile of the wadi (Schumm, 1973). This
29
30 body of sediments originating by the siltation mechanism may have been deposited relatively
31
32 quickly (**maximum a century**), considering sediment transport delivery rates of the wadi (~8500
33
34 m³/yr) (Achite and Ouillon, 2007) and the calculated reservoir volume of the dammed area
35
36 (~7.5*10⁵ m³). Maintenance and sweeping of the basin could certainly have delayed this process.
37
38
39
40
41
42
43
44
45
46
47
48
49

50 **The lower wadi**

51
52 Construction of the dam drastically reduced the runoff along the lower reach of the wadi,
53
54 which then became almost dry. We recognized a terrace that can be correlated only with the one
55
56 that pre-dates the construction of the dam (wadi terrace II, Fig. 1). Above these terraces two Roman
57
58 water cisterns were built, that should have supplied water to the town and probably to the Hadrianic
59
60 Baths (in construction in 123 A.D.) and, later, to the Severian Great *Nymphaeum* (Fig. 1). These

1
2
3 cisterns are located on the right bank of the wadi, fed by an aqueduct built by Q. Servilius Candidus
4
5 in 119-120 from Wadi Caam (20 km East), characterized by a more regular flow and ensuring a
6
7 water supply by connecting natural springs (i.e. connection aqueduct-Hadrianic Baths) (Romanelli,
8
9 1925; Bartoccini, 1929; Cifani and Munzi, 2003). Stratigraphical analysis in proximity of the upper
10
11 cistern (S4, Fig. 1) and on the left bank on top of the pre-dam terrace shows a deposit containing
12
13 Roman pottery and plaster fragments in a slightly compact silt. One piece of charcoal collected in
14
15 this layer yields a corrected age of 80-230 A.D. (Table I), implying that the construction of the dam
16
17 was completed before this time. Today the cisterns are partly covered by deposits derived from
18
19 siltation body as the wadi crosscut the dam infill and successively re-incised to reach a new
20
21 equilibrium.
22
23
24
25
26

27 The stratigraphy and morphology of Wadi Lebda provides evidence for subsequent phases
28
29 of extreme overflowing events on to the terraces and fan surfaces, followed by a return of the flow
30
31 inside the stream bed and formation of erosional terrace risers. Since the nearby Libyan coastline
32
33 can be considered tectonically stable (Florida, 1939; Lipparini, 1940; Capitanio, 1999; personal
34
35 observation of the Tyrrhenian (125ka) paleoshoreline), the discharge regimes, or liberation of
36
37 sediments, are likely not related to sea level changes. Consequently, the net stream longitudinal
38
39 profile adjustment produces only a limited downcutting that slightly deepens the channel floor.
40
41
42
43
44
45
46
47
48
49

50 **The harbor**

51 Nowadays, the 10^5 m² harbor basin is completely infilled by a depositional sequence that is
52
53 entrenched by the present wadi bed that here branches out into two channels. The infill surface
54
55 morphology gently deepens toward the sea, where it connects with the beach and the active sand
56
57 dunes located between the two dock tips (Fig. 4a). We studied the upper 4 meter of stratigraphy by
58
59 coring the sediments (S5, Fig. 1). Through sedimentological and micropaleontological analysis, we
60
recognized the occurrence of deposits from three different paleoenvironments that we describe

1
2
3 below from the bottom (~1.6 m below sea level) to the top (~2.2 m a.s.l.) of the sampled core (E1,
4
5 E2, E3, Fig. 4b):
6
7

8 E1) Gray sand with well-rounded quartz and a few small volcanic clasts, characterized by an
9
10 association of recent smoothed and well-preserved shallow water benthic foraminifera (e.g.
11
12 *Ammonia parkinsoniana*, *Peneroplis planatus*, *P. pertusus*, *Planorbulina mediterranensis*, *Sorites*
13
14 *orbicularis*, *Quinqueloculina* spp.). This indicates a marginal marine embayment, as expected in an
15
16 area such as the harbor before 210-400 A.D.;
17
18

19 E2) Gray fine sand with a strong decrease in well-preserved shallow water benthic
20
21 foraminifera (few specimens often smoothed). This indicates a protected paleoenvironment
22
23 probably not connected with the sea (the base of this layer date 210-400 A.D.; S5-2, Table I);
24
25
26

27 E3) Dark hazel fine sand and silt, upward graded, with weathered, well-rounded clasts and a
28
29 few smoothed benthic foraminifera, indicating major fluvial sediment transport (the base of this
30
31 layer date 560-660 A.D.; S5-1, Table I). The lower part of this latter deposit is anomalously
32
33 interrupted by a well-sorted quartzitic sand layer that could suggest the occurrence of an energetic
34
35 phase that affected the harbor. Toward the upper part, the deposit turns into reddish and orange sand
36
37 and fine sand, locally with small pebbles and bioturbations testifying to an alluvial and well-defined
38
39 continental environment.
40
41
42

43 Previous authors have suggested that the harbor was never used because it was shortly filled
44
45 with sediments (Salza Prina Ricotti, 1972-1973). However, our data suggest that the harbor basin
46
47 hosted marine water (at least before 210-400 A.D.) and thus we suggest that the harbor built by
48
49 Septimius Severus was functional. Then, its marine sedimentation progressively changed and the
50
51 leading depositional process became that related to alluviation episodes due to sediment delivery
52
53 from the Wadi Lebda. The coring location, close to the eastern docks, is the part of the harbor most
54
55 prone to infilling and thus has the highest probability for sediment preservation. In the western part
56
57 of the harbor, where the wadi flows nowadays, the alluvial infill was limited. Consequently, this
58
59 part of the harbor basin was probably used for a longer time. This hypothesis is also supported by
60

1
2
3 the fact that a productive/residential settlement was sited at the beginning of the V century between
4 the western docks and the western beach and survived for all the Byzantine period. During Arabic
5 times the settlement also expanded on to the alluvial deposits within the harbor basin (Laronde,
6 1988), suggesting a landscape very comparable with that of today and at the end of harbor use.
7
8
9
10
11
12
13
14
15
16

17 Conclusions

18
19 Figure 5 shows how we correlated the data of the different sites of investigation in order to
20 reconstruct the wadi evolution/response in time.
21
22
23

24 Wadi Lebda was responsible during the Holocene for the formation of an alluvial plain upon
25 which Leptis Magna was built (A, Fig. 6). Still, it plays a dominant role in the production, transport
26 and deposition of detritus into both the town and the harbor as sedimentary basins by flash floods.
27
28
29
30

31 In order to mitigate the severe flooding risk that was threatening the settlement, the Romans
32 constructed the dam and excavated the ditch along the *agger*. Both the dam and the *agger* acted to
33 control the Wadi Lebda and protect the town and harbor from possible overfloodings by decreasing
34 the energy and draining out the intermittent flow from the wadi (B, Fig. 6).
35
36
37
38
39
40

41 Construction of the dam modified the behavior of the wadi, inducing longitudinal profile
42 adjustment with sediment overloading and consequent aggradation and siltation in the artificial
43 reservoir. Conversely, the lower reach of the wadi began to have a degradational behavior,
44 preventing the excess of sediment delivery into the harbor basin and favoring marine deposition, as
45 promoted by: (1) late sediment-deficient alluvial discharge; (2) predominant canalized runoff.
46 According to statistical analysis of the sequence of ages of the analyzed deposits ($S_3 < S_2 < S_4$, i.e.
47 pre-dam deposits younger than the *agger*/dam construction, in turn younger than post-dam
48 deposits), the *agger*/dam system was built between 32-131 A.D. (considering the 95.4 % of
49 probability) (Table I). This is in agreement with historical and archaeological literature that identify
50 two events related to the *agger*/dam system: the defensive need in 69 A.D. against the Garamantes
51
52
53
54
55
56
57
58
59
60

1
2
3 (Tacitus, 100 A. D.; Goodchild and Ward Perkins, 1949) and the building of the hydraulic network
4
5 in 119-123 (aqueduct from Wadi Caam and Hadrianic Baths) occupying part of the dried wadi bed
6
7
8 (C, Fig. 6).
9

10 After a period in which the agger/dam system was fully functional, extending at least up to
11
12 the period of construction and use of the monumental harbor (beginning of III century), the
13
14 **maintenance of the system was stopped and siltation quickly filled the reservoir.** When siltation
15
16 reached the top of the dam the wadi was free to flow on this upper surface and, spilling out from the
17
18 dam, by-passed the construction and started flowing again into the lower wadi reach, producing
19
20 limited alluviation episodes in the abandoned wadi bed and at the harbor (D, Fig. 6). Because the
21
22 reservoir is expected to completely fill within 100 yrs maximum, most likely before the beginning
23
24 of the harbor alluviation (210-400 A.D. from the stratigraphy of the cores, Fig. 5), maintenance of
25
26 the agger/dam system was probably stopped at some point between the III and beginning of IV cent.
27
28
29
30

31 **Water spilling out of the dam likely favored erosion at the base of the dam** itself by affecting
32
33 the stability of its right shoulder, resulting in complete collapse. At this point the wadi gained again
34
35 its freedom to erode and flood wide areas originally reclaimed by the Lepticians. In fact, rupture of
36
37 the dam produced a double effect on the wadi longitudinal profile adjustment: erosion of the
38
39 siltation deposits and strong remobilization of sediments, with sediment overloading from the dam
40
41 point source resulting in channel-bed aggradation (i.e. the downfilling (Schumm, 1993; Sugai,
42
43 1993)) and **consequent harbor infill** (E, Fig. 6). The Town was buried under alluvial deposits by
44
45 consecutive flooding events after 320-440 A.D., thus, the dam rupture did not occur later than the
46
47 IV- beginning of V cent. (Fig. 5).
48
49
50
51

52 **The structure of the dam does not show any type of damage that could have been caused by**
53
54 **a local earthquake or by the occurrence of a destructive earthquake that hit a large part of the**
55
56 **eastern Mediterranean, such as the 365 A.D. Creta earthquake.** Moreover, the fact that the dam
57
58 basin is filled by siltation sediments supports the hypothesis that the cause of the damage is related
59
60

1
2
3 to lack of maintenance and not to an unexpected event such as an earthquake. The latter would have
4
5 hit the dam in functioning conditions.
6
7

8 **Lack of maintenance and rupture of the dam signaled the beginning of the decline of the**
9
10 **harbor. In fact, as deduced by the core stratigraphy and dating, the harbor basin started first to be**
11 **infilled by limited alluvial deposits between 210 and 400 A.D.** (D, Fig. 6), then it turned into a
12
13 purely continental environment before 560-660 A.D. (E, Fig. 6).
14
15
16

17 **No evidence for a clear event of tsunami** inundation hitting the harbor, or for an important
18
19 change in the wadi-sea level balance during the past couple of millennia (F, Fig. 6) were found (the
20
21 present bottom of the wadi in the harbor area is only a few centimeters below the one pre-dating the
22
23 dam construction).
24
25
26

27 The change in the balance between natural disasters and the vulnerability of society,
28
29 according to its socio-economic state, make the correct siting of settlements and infrastructures
30
31 time-dependent. Until the Romans had the economic possibility to face the siltation troubles within
32
33 the dam reservoir, by ensuring the maintenance of the hydraulic system, the town was protected.
34
35 But, as the resources to counteract natural processes become less efficient and less widely available,
36
37 nature once again began to dominate. A society that today appears to be resilient to natural
38
39 perturbations can often prove unable to cope in the future. On this note, as “deciphering the past is a
40
41 key to the future” (Charles, 1873), we must be aware that since we have urbanized areas that are
42
43 known to be vulnerable to repeated and extreme natural events, we need to shield them, regardless
44
45 of the economic fluctuations that tend to change societal priorities.
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Acknowledgements

We are grateful to Prof. R. Funiciello for his enthusiastic support in motivating and inspiring this research, and to F. Baroni and F. Felici for the constructive sustain and comments.

We thank the participants to the Italian Archeological Mission of the University of Roma³ in Leptis Magna led by L. Musso: L. Marsico, B. Bianchi, L. Buccino and C. Malaguti.

A special acknowledge to the Department of Archeology of Libya, in particular to Dr. M. Massaud of Department of Antiquities of Leptis Magna.

For Peer Review

References

Achite, M, and Ouillon, S. 2007: Suspended sediment transport in a semiarid watershed, Wadi Abd, Algeria (1973-1995). *Journal of Hydrology*, 343, 187-202.

Ambraseys, N. N., Melville, C. P., and Adams, R. D. 1994: *The Seismicity of Egypt, Arabia and the Red Sea: A Historical Review*, Cambridge Univ. Press, Cambridge.

Bartoccini, R. 1929:, *Le terme di Lepcis (Leptis Magna)*, Istituto italiano d'arti grafiche, Bergamo, 203 pp.

Bronk Ramsey, C. 2009: Bayesian analysis of radiocarbon dates. *Radiocarbon*, 51(1), 337-360.

Capitanio, F. A. 1999: Aspetti della tettonica terziaria in Tripolitania, Libia, M.S. Thesis, University of Roma 3.

Charles, L. 1873:, *The Geological Evidences of The Antiquity of Man*, (4th ed.), New York: AMS Press.

Cifani, G., and Munzi, M. 2003: Alle sorgenti del Cinyps. *Libyan Studies*, 34, 85-100.

Di Vita, A. 1990: Sismi, urbanistica e cronologia assoluta. Terremoti e urbanistica nella città di Tripolitania tra il I secolo a.C. ed il IV d.C. in: *L'Afrique dans L'Occident Romani (Ier Siècle Av. J.-C.- IVe Siècle Ap. J.- C.) Actes du Colloqui Organisé par l'Èrcole Français de Rome sous le Patronage de l'Institut National d'Archéologie et d'Art de Tunis (Rome, 3-5 décembre 1987)*, Roma, 425-494.

Di Vita, A. 1995: Archaeologists and earthquakes: the case of 365 A.D. *Annali di Geofisica*, 38, 971-976.

1
2
3
4
5 Florida, G. B. 1939: Osservazioni sul Miocene dei dintorni di Homs. *Boll. Soc. Geol. It. LVIII*.

6
7
8
9
10 Goodchild, R. G., and Ward Perkins J. B. 1949: The Roman and Byzantine defenses of Lepcis
11 Magna. *PBSR*, 21, 42-73.

12
13
14
15
16 Goslar, T., Czernik, J., and Goslar, E. 2004: Low-energy ¹⁴C AMS in Poznan' Radiocarbon
17 Laboratory, *Nuclear Instruments and Methods. B*, 223-224, 5-11.

18
19
20
21
22
23 Guidoboni, E., Comastri, A. and Traina, G. 1994: *Catalogue of ancient earthquakes in the*
24 *Mediterranean area up to the 10th century*, SGA Editor, Bologna. ING, Rome.

25
26
27
28
29 Laronde, A. 1988: Le port de Lepcis Magna. *Comptes-rendus des séances de l'Académie des*
30 *inscriptions et belles-lettres*, 132 (2), 337-353.

31
32
33
34
35 Lipparini, T. 1940: Tettonica e geomorfologia della Tripolitania. *Boll. Soc. Geol. It.*, LIX.

36
37
38
39
40 Lorito, S., Tiberti, M. M., Basili, R., Piatanesi, A. and Valensise, G. 2007: Earthquake-generated
41 tsunamis in the Mediterranean Sea: scenarios of potential threats to Southern Italy. *J. Geophys. Res.*, doi:
42 10.1029/2007JB004943.
43
44
45

46
47
48
49 Reid, I., and Laronne, J. 1995: Bedload sediment transport in an ephemeral stream and a comparison
50 with seasonal and perennial counterparts. *Water Resources Research*, 31 (3), 773–781.

51
52
53
54
55 Reimer, P. J., L. Baillie, M. G., Bard, E., Bayliss, A., Beck, J. W., Bertrand, C. J. H., Blackwell, P.
56 G., Buck, C. E., Burr, G. S., Cutler, K. B., Damon, P. E., Edwards, R. L., Fairbanks, R. G., Friedrich, M.,
57 Guilderson, T. P., Hogg, A. G., Hughen, K. A., M. Kromer, B. G., Manning, S., Ramsey, C. B., Reimer, R.
58 W., Remmele, S., Southon, J. R., Stuiver, M., Talamo, S., Taylor, F. W., van der Plicht, J., and
59
60

1
2
3 Weyhenmeyer, C. E. 2004: IntCal04-terrestrial radiocarbon age calibration, 0--26 cal kyr B. *Radiocarbon*, 46
4
5 (3), 1029-1058.
6
7

8
9 Rinaldi Tufi, S. 2000: *Archeologia delle province romane*, Carocci Ed., Roma, 222 pp.
10
11

12
13 Romanelli, P. 1925: *Leptis Magna, Africa Italiana*, Roma.
14
15

16
17 Sallust 40 B.C.: *Bellum Iugurthinum*, 77, 1-3.
18
19

20
21 Salza Prina Ricotti, E. 1972-1973: I porti della zona di Leptis Magna. *Rendiconti della Pontificia*
22
23 *Accademia di Archeologia*, 45, 75-103.
24
25

26
27 Salza Prina Ricotti, E. 1995: Lepis Magna la città delle ombre bianche. *Archeo*, 9 (127), 50-91.
28
29

30
31 Schick, A. P. 1977: Study of sediment generation, transport and deposition in semiarid zones.
32
33 *Hydrol. Sci. Bull.*, 22 (4), 535-542.
34
35

36
37 Schumm, S. A. 1973: Geomorphic thresholds and complex response of drainage systems. In *Fluvial*
38
39 *Geomorphology*, M. Morisawa Editor, Binghamton State University, Binghamton, NY, 299-310.
40
41

42
43 Schumm, S. A. 1993: River response to baselevel change; implications for sequence stratigraphy.
44
45 *Journal of Geology*, 101 (2), 279-294.
46
47

48
49 Serrat, P., Ludwig, W., Navarro, B., and Blazi, J.L. 2001: Spatial and temporal variability of
50
51 sediment fluxes from a coastal Mediterranean river: the Têt (France). *Comptes Rendus de l'Académie des*
52
53 *Sciences IIA*, 333 (7), 389-397.
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Shaw, B., Ambraseys, N. N., England, P. C., Floyd, M. A., Gorman, G. J., Higham, T. F. G., Jackson, J. A., Nocquet, J.-M., Pain, C. C. and Piggott, M. D. 2008: Eastern Mediterranean tectonics and tsunami hazard inferred from the AD 365 earthquake. *Nature Geoscience*, 1, 268-276, doi:10.1038/ngeo151.

Stiros, S. C. 2001: The AD 365 Crete earthquake and possible seismic clustering during the fourth to sixth centuries AD in the Eastern Mediterranean: a review of historical and archaeological data. *Journal of Structural Geology*, 23, 545-562.

Stuiver, M., and Reimer, P. J. 1993: Extended (super 14) C data base and revised CALIB 3.0 (super 14) C age calibration program. *Radiocarbon*, 35, 215-230.

Sugai, T. 1993: River terrace development by concurrent fluvial processes and climate change. *Geomorphology*, 6, 243-252.

Tacitus 100 A. D.: *Historiae*, 4.50.

Captions

Table I.

Radiocarbon ages and associated chronological data for sediments from the Leptis Magna area. The calibrated ages are calculated for the probability density function at the 95.4% (2σ) and at 68.2% (1σ) confidence limits. *Statistical analysis of sequential dates by means of: OxCal v4.1 © by Bronk Ramsey (2009); IntCal04 atmospheric curve (Reimer *et al.*, 2004).

Figure 1. Geological and geomorphological map. The correlated wuadi terraces (I-III) and the stratigraphic sections (S1-S5) are reported. The inset shows the location of the study area.

Figure 2. Alluvial deposits of the town. a) the oldest polycyclic deposits that buried the roman buildings; b) alluvial deposits dip; c) view of the alluvial fan surface overlaid by sand dunes.

Figure 3. The Roman Dam. a) frontal view of the dam. On the left is visible the pre-dam wadi bed; b) view of the terrace formed by the dam's reservoir siltation; c) Stratigraphic section of the reservoir infill. Both deposits, laminated for the wadi flow and massive for siltation of the reservoir, in the lower and upper parts, respectively, are evident.

Figure 4. The harbor. a) view of the infilled harbor basin. b) Stratigraphic column of the cored deposits. Location of samples for ^{14}C (white square), micropaleontological and mineralogical (gray square) analyses is reported along with photos of ceramics and FESEM view of sand and some selected benthic foraminifera. The FESEM picture 1 shows not well sorted sand with dominant, smoothed quartz. The FESEM pictures of some benthic foraminifera (scale bar represents 100 μm) show: 2) *Ammonia parkinsoniana* and 3) *Peneroplis planatus*.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure 5. Correlation of the results of the study sites. The overlap between consecutive period of events is related only to 2σ confidence limits.

Figure 6. Conceptual sketch of the evolution of the Wadi Lebda longitudinal profile and its depositional history.

For Peer Review

ID	Radiocarbon age (yr BP)	% area enclosed	Cal. A.D. age ranges (yr A.D.)	Rel. area under prob. distrib. [Reimer et al., 2004]	Applied age ranges (yr A.D.)
S1	1655 ±35	68.3 (1σ)	344-427	1.000	340-430
		95.4 (2σ)	259-285	0.056	320-440
			288-292	0.005	
			322-443	0.826	
			448-463	0.017	
			483-532	0.096	
S2	1920 ± 35	68.3 (1σ)	54-126	1.000	50-130
		95.4 (2σ)	1-143	0.954	0-140
			146-173	0.027	
			193-210	0.019	
S3	1960± 30	68.3 (1σ)	5-12	0.091	20-70
			16-72	0.909	
		95.4 (2σ)	39 B.C.-87 A.D.	0.961	40 B.C.-90 A.D.
			104-121	0.039	
S4	1865± 30	68.3 (1σ)	86- 107	0.222	120-180
			120-175	0.579	
			191-211	0.199	
		95.4 (2σ)	77-228	1.000	80-230
S5	1435± 35	68.3 (1σ)	601-648	1.000	600-650
		95.4 (2σ)	564-658	1.000	560-660
	1745±35	68.3 (1σ)	244-337	1.000	240-340
		95.4 (2σ)	181-185	0.003	210-400
			214-401	0.997	
S3>S2>S4*	1920 ± 35	68.3 (1σ)	59-96	0.846	30-130
			104-115	0.154	
		95.4 (2σ)	32-131	1.000	

Table I.

Radiocarbon ages and associated chronological data for sediments from the Leptis Magna area. The calibrated ages are calculated for the probability density function at the 95.4% (2σ) and at 68.2% (1σ) confidence limits. *Statistical analysis of sequential dates by means of: OxCal v4.1 © by Bronk Ramsey (2009); IntCal04 atmospheric curve (Reimer *et al*, 2004).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Fig. 1

169x208mm (300 x 300 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Fig. 2

263x119mm (300 x 300 DPI)

Peer Review

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



Fig. 3

229x138mm (300 x 300 DPI)

Review

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Fig. 4

168x189mm (300 x 300 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Fig.5

176x146mm (300 x 300 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Fig.6

136x214mm (300 x 300 DPI)